

Structural properties of acyclic heaps with applications to Kazhdan–Lusztig theory

Brent G. Pohlmann

Ph.D., University of Colorado at Boulder, 2008

October 5, 2008

Abstract

Structural properties of acyclic heaps of pieces with Kazhdan–Lusztig theory

Thesis directed by Prof. Richard M. Green

Let W be an arbitrary Coxeter group with generating set S of involutions. A reduced expression for an element $w \in W$ is a minimal length word in S that represents w . The set W_c of fully commutative elements are characterized by the property that any reduced expression for $w \in W_c$ can be obtained from any other via iterated commutations of adjacent generators. If $w \in W_c$ and $sw \notin W_c$ for some $s \in S$, then we say sw is weakly complex.

Star reducible Coxeter groups are a class of Coxeter groups whose fully commutative elements have a particularly nice property. Star reducible Coxeter groups contain the finite Coxeter groups as a subclass.

A heap is an isomorphism class of labelled posets. Each heap is equipped with a set of edges and a set of vertices. Green defined a linear map ∂_E which sends each edge of a heap E to a linear combination of vertices. If $v \in \text{Im} \partial_E$, we call v a boundary vertex. If e_0 is an edge of E and $\partial_E(e_0) = v$ for a vertex v , then we call v an effective boundary vertex. If $\partial_E(e_0) = v_1 + v_2$, then v_1 and v_2 are said to be linearly equivalent. A result by Stembridge states that every fully commutative element has a unique heap.

The main result we will prove here is that in the heap of a fully commutative element in a star reducible Coxeter group, every boundary vertex is linearly equivalent to an effective boundary vertex.

We use the main result to prove another theorem concerning the basis elements $\tilde{t}_w : w \in W_c$ of a generalized Temperley–Lieb algebra. The theorem we prove allows an inductive computation of the c -basis which is a linear combination of the $\tilde{t}_w : w \in W_c$. Furthermore, the c -basis can be shown to have nonnegative structure constants, that is, structure constants that are Laurent polynomials with nonnegative coefficients. One of the reasons this is interesting is that in many cases, these structure constants are also structure constants for the well known Kazhdan–Lusztig basis C'_w , whose positivity is generally very difficult to prove.

Contents

1	Introduction	1
2	Coxeter Group Theory	3
2.1	Introduction	3
3	Heaps of Pieces	9
3.1	Introduction	9
3.2	The Boundary Map ∂	11
3.3	Some Properties of Heaps and Full Commutativity	12
3.4	Main Theorem	17
4	Property W	20
4.1	Introduction	20
4.2	Acyclic Case	21
4.3	Forbidden Configurations	23
4.4	The non-acyclic case	26

Chapter 1

Introduction

A Coxeter group W is a group with finite generating set S and presentation given by

$$W = \langle S \mid (st)^{m(s,t)} = 1 \text{ for } m(s,t) < \infty \rangle.$$

Each $w \neq 1$ in W can be written in the form $w = s_1 s_2 \cdots s_r$ for some $s_i \in S$. If r is as small as possible we call r the length of w , written $\ell(w)$. A reduced expression for an element $w \in W$ is a minimal length word in S that represents w . The set W_c of fully commutative elements of W is characterized by the property that any reduced word for $w \in W_c$ can be obtained from any other via iterated applications of short braid relations, that is, relations of the form $st = ts$, where $s, t \in S$. For example, if w is a product of commuting generators from S , then w is fully commutative. If $w \in W_c$ and $sw \notin W_c$ for some $s \in S$, then we say sw is weakly complex. Green [6] defined the star reducible Coxeter groups to be those Coxeter groups for which every fully commutative element is equivalent to a product of commuting generators by a sequence of length-decreasing star operations. The star reducible Coxeter groups are defined and listed in Section 2.1.

Denote by $\mathcal{H}_q = \mathcal{H}_q(X)$ the Hecke algebra associated to W . This is a $\mathbb{Z}[q, q^{-1}]$ -algebra with a basis consisting of (invertible) elements $\{T_w : w \in W\}$, and an associative multiplication. For our purposes, we extend the scalars of $\mathcal{H}_q$ by setting $v^2 = q$ and writing $\mathcal{H} = \mathcal{A} \otimes_{\mathbb{Z}[q, q^{-1}]} \mathcal{H}_q$ where $\mathcal{A} = \mathbb{Z}[v, v^{-1}]$. We write $\mathcal{A}^+$ and $\mathcal{A}^-$ for $\mathbb{Z}[v]$ and $\mathbb{Z}[v^{-1}]$, respectively. We also define a scaled version of the T -basis, $\{\tilde{T}_w : w \in W\}$, where $\tilde{T}_w := v^{-\ell(w)} T_w$. In [10], Kazhdan and Lusztig defined the bases $\{C_w : w \in W\}$ and $\{C'_w : w \in W\}$ for $\mathcal{H}$. These Kazhdan–Lusztig bases are constructed from the bases $\{T_w : w \in W\}$. An equation relating the C'_w -basis and the $\tilde{T}_w$ -basis is given by

$$C'_w = \tilde{T}_w + \sum_{y < w} A_{y,w} \tilde{T}_y$$

where $<$ is the Bruhat order on W and $A_{y,w} \in v^{-1}\mathcal{A}^-$. Following [3, Section 11.1], we denote the coefficients of $\tilde{T}_y$ in C'_w by $P_{y,w}^*(q)$. The Kazhdan–Lusztig polynomial $P_{y,w}$ is then given by $v^{\ell(w)-\ell(y)} P_{y,w}^*(q)$. The coefficient of the v^{-1} term in $P_{y,w}^*(q)$ is $\mu(y, w)$, which is very difficult to compute efficiently, even for moderately small groups.

Let $J(X)$ be the two-sided ideal of $\mathcal{H}$ generated by the elements

$$\sum_{w \in \langle s, t \rangle} T_w,$$

where (s, t) runs over all pairs of elements of S such that $2 < m(s, t) < \infty$ and $\langle s, t \rangle$ is the parabolic subgroup generated by s and t . Following Graham [4], we define the generalized Temperley–Lieb algebra $TL(X)$ to be the quotient $\mathcal{A}$ -algebra $\mathcal{H}(X)/J(X)$ and denote the corresponding epimorphism of algebras by $\theta : \mathcal{H}(X) \rightarrow TL(X)$. For $\tilde{T}_w \in \mathcal{H}$, we have $\theta(\tilde{T}_w) = \tilde{t}_w$ and define the $\mathcal{A}^-$ -submodule L of $TL(X)$ to be that generated by the $\{\tilde{t}_w : w \in W_c\}$.

Green and Losonczy [8] proved that for each $w \in W_c$ there exists a unique $c_w \in TL(X)$ such that

$$c_w = \tilde{t}_w + \sum_{\substack{y < w \\ y \in W_c}} a_{y,w} \tilde{t}_y,$$

where $<$ is the Bruhat order on W , and $a_{y,w} \in \mathcal{A}^-$ for all y . We note here that the $c_w \in TL(X)$ are analogous to $C'_w \in \mathcal{H}$ and the direct connection, which we explore later, is not obvious.

A heap is an isomorphism class of labelled posets satisfying certain axioms. Each heap is equipped with a set of edges and a set of vertices. In [5], Green defined a linear map ∂_E which sends each edge of E to a linear combination of vertices. We say v is a boundary vertex if $v \in \text{Im } \partial_E$ and that v is an effective boundary vertex if $\partial_E(e_1) = v$ for an edge e_1 from E . Two boundary vertices v_1 and v_2 are said to be linearly equivalent if $\partial_E(e_1) = v_1 + v_2$ for some edge e_1 from E . In §3.4 we prove that in the heap of a fully commutative element in a star reducible Coxeter group, every boundary vertex is linearly equivalent to an effective boundary vertex. This is a subtle structural property of the heap and the proof is combinatorial in nature.

As an application of Theorem 3.4.1, we prove that if $x \in W$ is weakly complex for W star reducible, we have $\tilde{t}_x \in v^{-1}\mathcal{L}$. This property of the $\tilde{t}_x$, which Green calls Property W in [7], allows the inductive computation of the c -basis using the formula

$$c_s c_w = \begin{cases} (v + v^{-1})c_w, & \text{if } \ell(sw) < \ell(w), \\ c_{sw} + \sum_{sy < y} \mu(y, w) c_y & \text{if } \ell(sw) > \ell(w). \end{cases}$$

Under Property W, the c -basis can be shown to have nonnegative structure constants, that is, structure constants that are Laurent polynomials with nonnegative coefficients. One of the reasons this is interesting is that in many cases, these structure constants are also structure constants for the Kazhdan–Lusztig basis C'_w , whose positivity is generally very difficult to prove.

Chapter 2

Coxeter Group Theory

2.1 Introduction

In §2 we review general Coxeter group theory. For a complete review of this material, the reader is referred to [9] and [2].

A Coxeter system (W, S) consists of a group W with distinguished (finite) set of generating involutions S and presentation given by

$$W = \langle S \mid (st)^{m(s,t)} = 1 \text{ for } m(s,t) < \infty \rangle,$$

where $m(s, s) = 1$. (It turns out that the $s \in S$ are distinct as group elements, and that $m(s, t)$ is the order of st .) Since the generators $s \in S$ have order 2 in W , each $w \neq 1$ in W can be written in the form $w = s_1 s_2 \cdots s_r$ for some $s_i \in S$. If r is as small as possible we call r the *length* of w , written $\ell(w)$. A product $w_1 w_2 \cdots w_n$ of elements $w_i \in W$ is called *reduced* if $\ell(w_1 w_2 \cdots w_n) = \sum_i \ell(w_i)$. We reserve the terminology *reduced expression* for reduced products $w_1 w_2 \cdots w_n$ in which every $w_i \in S$. To specify a Coxeter system (W, S) we can specify a finite set S and draw an undirected graph X with set S as vertex set, joining vertices s and t by an edge labelled $m(s, t)$ whenever this number is at least 3. If distinct vertices s and t are not joined, it is understood that $m(s, t) = 2$. As a simplifying convention, the label $m(s, t) = 3$ is usually omitted. We call X a *Coxeter graph*. The connected graph in Figure 1 is called a Coxeter graph of type B_4 .




FIG. 1. Coxeter graph of type B_4 .

We call an element $w \in W$ *complex* if it can be written as a reduced product $x_1 w_{st} x_2$, where $x_1, x_2 \in W$ and w_{st} is the longest element of some rank 2 parabolic subgroup $\langle s, t \rangle$ such that s and t correspond to adjacent nodes in the Coxeter graph. An element $w \in W$ is said to be *weakly complex* if (a) it is complex and (b) it is of the form $w = su$, where $s \in S$ and u is not complex. We write

$$\mathcal{L}(w) = \{s \in S : \ell(sw) < \ell(w)\}$$

and

$$\mathcal{R}(w) = \{s \in S : \ell(ws) < \ell(w)\}.$$

The set $\mathcal{L}(w)$ (respectively, $\mathcal{R}(w)$) is called the *left* (respectively, *right*) *descent set* of w .

Example 2.1.1 *The reduced element $w = s_1 s_2 s_1 s_2 s_3$ in type B_4 is complex: the subexpression $w' = s_1 s_2 s_1 s_2$ is the longest element of the rank 2 parabolic subgroup $\langle s_1, s_2 \rangle$. If we delete the first occurrence of s_1 in w we are left with an element that is not complex. Thus w is also weakly complex, with $s = s_1$ in the definition above.*

Denote by $W_c(X)$ the set of all elements of W that are not complex. The elements of $W = W_c(X)$ are the *fully commutative* elements of [12]; they are characterized by the property that any two of their reduced expressions may be obtained from each other by repeated commutation of adjacent generators.

Denote by $\mathcal{H}_q = \mathcal{H}_q(X)$ the Hecke algebra associated to W . This is a $\mathcal{A}$ -algebra with a basis consisting of (invertible) elements $T_w : w \in W$, satisfying

$$T_s T_w = \begin{cases} T_{sw}, & \text{if } \ell(sw) > \ell(w) \\ qT_{sw} + (q-1)T_w, & \text{if } \ell(sw) < \ell(w) \end{cases}$$

where ℓ is the length function on the Coxeter group W , $w \in W$, and $s \in S$. Note that for $s \in S$, we have $T_s^2 = (q-1)T_s + qT_1$.

If $w = s_1 s_2 \cdots s_r$ is a reduced expression for w , then $T_w = T_{s_1} T_{s_2} \cdots T_{s_r}$. Since $\mathcal{H}$ can also be shown to be associative, the product $T_x T_{x'}$ is defined for all $x, x' \in W$.

Using the above relations we can show that for all $s \in S$:

$$T_s^{-1} = q^{-1}T_s - (1 - q^{-1})T_1.$$

Therefore every T_w is invertible in $\mathcal{H}$.

For our purposes, we extend the scalars of $\mathcal{H}_q$ by setting $v^2 = q$ and define $\mathcal{H} = \mathcal{A} \otimes_{\mathbb{Z}[q, q^{-1}]} \mathcal{H}_q$. We also define a scaled version of the T -basis, $\{\tilde{T}_w : w \in W\}$, where $\tilde{T}_w := v^{-l(w)} T_w$. We will write $\mathcal{A}^+$ and $\mathcal{A}^-$ for $\mathbb{Z}[v]$ and $\mathbb{Z}[v^{-1}]$, respectively. We denote the $\mathbb{Z}$ -linear ring homomorphism $\mathcal{A} \rightarrow \mathcal{A}$ exchanging v and v^{-1} by $\bar{}$. We can extend $\bar{}$ to a ring automorphism of $\mathcal{H}$ by the condition that

$$\overline{\sum_{w \in W} a_w T_w} = \sum_{w \in W} \bar{a}_w T_w^{-1}$$

where the a_w are elements of $\mathcal{A}$.

Let $J(X)$ be the two-sided ideal of $\mathcal{H}$ generated by the elements

$$\sum_{w \in \langle s, t \rangle} T_w,$$

where (s, t) runs over all pairs of elements of S that correspond to adjacent nodes in the Coxeter graph, and $\langle s, t \rangle$ is the (finite) parabolic subgroup generated by s and t . Following Graham [4], we define the *generalized Temperley–Lieb algebra* $TL(X)$ to be the quotient $\mathcal{A}$ -algebra $\mathcal{H}(X)/J(X)$. We denote the corresponding epimorphism of algebras by $\theta : \mathcal{H}(X) \rightarrow TL(X)$. Let t_w (respectively, $\tilde{t}_w$) denote the image in $TL(X)$ of the basis element T_w (respectively, $\tilde{T}_w$) of $\mathcal{H}$.

In [10], Kazhdan and Lusztig defined the bases $\{C_w : w \in W\}$ and $\{C'_w : w \in W\}$ for $\mathcal{H}$. These Kazhdan–Lusztig bases are constructed from the bases $\{T_w : w \in W\}$. The following theorem can be viewed as a restatement of [10, 1.1.c].

Theorem 2.1.2 (Kazhdan, Lusztig) *For each $w \in W$, there exists a unique $C'_w \in \mathcal{H}$ such that both $\overline{C'_w} = C'_w$ and*

$$C'_w = \tilde{T}_w + \sum_{y < w} A_{y,w} \tilde{T}_y,$$

where $<$ is the Bruhat order on W and $A_{y,w} \in v^{-1}\mathcal{A}^-$ are certain polynomials in v^{-1} .

The basis $\{C_w\}_{w \in W}$ is closely related to $\{C'_w\}_{w \in W}$, and it suffices to understand one of them.

We define the $\mathcal{A}^-$ submodule $\mathcal{L}$ of $TL(X)$ to be that generated by $\{\tilde{t}_w : w \in W_c\}$. We define $\pi : \mathbb{L} \rightarrow \mathbb{L}/v^{-1}\mathbb{L}$ to be the canonical $\mathbb{Z}$ -linear projection. It was proved in [8, Lemma 1.4] that the ideal $J(X)$ is fixed by $\bar{}$, so $\bar{}$ induces an involution on $TL(X)$ that sends v to v^{-1} and t_w to $t_{w^{-1}}^{-1}$. We denote this map also by $\bar{}$.

The following theorem establishes a canonical basis for $TL(X)$ in terms of the $\tilde{t}$ -basis. It comes from [8].

Theorem 2.1.3 *For each $w \in W_c$, there exists a unique $c_w \in TL(X)$ such that both $\overline{c_w} = c_w$ and $\pi(c_w) = \pi(\tilde{t}_w)$. Furthermore, we have*

$$c_w = \tilde{t}_w + \sum_{\substack{y < w \\ y \in W_c}} a_{y,w} \tilde{t}_y,$$

where $<$ is the Bruhat order on W , and $a_{y,w} \in v^{-1}\mathcal{A}^-$ for all y .

For later purposes, we define the following sublattices of the $\mathcal{A}^-$ -lattice $\mathcal{L}$.

Definition 2.1.4 *Let $s \in S$. We define $\mathcal{L}_L^s$ to be the free $\mathcal{A}^-$ -module with basis*

$$\{\tilde{t}_w : w \in W_c, sw < w\} \cup \{v^{-1}\tilde{t}_w : w \in W_c, sw > w\}.$$

Similarly, we define a free $\mathcal{A}^-$ -module $\mathcal{L}_R^s$.

Star operations are of key importance to this thesis. These were introduced in the simply laced case in [10] and in general in [11].

Definition 2.1.5 *Let $W = W(X)$ be any Coxeter group and let $I = \{s, t\} \subseteq S$ be a pair of noncommuting generators whose product has order $m = m(s, t)$ (where ∞ is allowed). Let W^I denote the set of all $w \in W$ satisfying $\mathcal{L}(w) \cap I = \emptyset$. Standard properties of Coxeter groups [9] show that any element $w \in W$ may be uniquely written as $w = w_I w^I$, where $w_I \in W_I = \langle s, t \rangle$, $w^I \in W^I$ and $\ell(w) = \ell(w_I) + \ell(w^I)$. There are four possibilities for the elements $w \in W$:*

- (i) *w is the shortest element in the coset $W_I w$, so $w_I = 1$ and $w \in W^I$;*
- (ii) *w is the longest element in the coset $W_I w$, so w_I is the longest element of W_I (which can only happen if W_I is finite);*
- (iii) *w is one of the $(m-1)$ elements $sw^I, tsw^I, stsw^I, \dots$;*
- (iv) *w is one of the $(m-1)$ elements $tw^I, stw^I, tstw^I, \dots$.*

The sequences appearing in (iii) and (iv) are called (left) $\{s, t\}$ -strings, or *strings* if the context is clear. If x and y are two elements of an $\{s, t\}$ -string such that $\ell(x) = \ell(y) - 1$, we call the pair $\{x, y\}$ *left $\{s, t\}$ -adjacent*, and we say that y is *left star reducible* to x . The above concepts all have

right-handed counterparts, leading to the notion of *right* $\{s, t\}$ -adjacent and *right star reducible* pairs of elements, and coset decompositions $({}^I w)_I$. If there is a (possibly trivial) sequence

$$x = w_0, w_1, \dots, w_k = y$$


where, for each $0 \leq i < k$, w_{i+1} is left star reducible or right star reducible to w_i with respect to some pair $\{s, t\}$, we say that y is *star reducible* to x . Because star reducibility decreases length, this relation is antisymmetric and thus defines a partial order on W . If w is an element of a left $\{s, t\}$ -string, S_w , we have $\{\ell(sw), \ell(tw)\} = \{\ell(w) - 1, \ell(w) + 1\}$; let us assume without loss of generality that sw is longer than w and tw is shorter. If sw is an element of S_w , we define $*w = sw$; if not, $*w$ is undefined. If tw is an element of S_w , we define ${}_w w = tw$; if not, ${}_w w$ is undefined. There are also obvious right handed analogues to the above concepts, so the symbols w^* and w_* may be used with the analogous meanings.

Example 2.1.6 Let W the Coxeter group of type B_4 and let $I = \{s_1, s_2\}$, where $m(s_1, s_2) = 4$. If we consider the element $w = s_2 s_1$, we have

$${}_w w = s_1, {}^* w = s_1 s_2 s_1, w_* = s_2 \text{ and } w^* = s_2 s_1 s_2.$$

If $x = s_1 s_2 s_1$ then $*x$ is undefined and x^* is undefined; if $x = s_2$ then $*x$ and x_* are undefined.

Definition 2.1.7 We say that a Coxeter group $W(X)$, or its Coxeter graph X , is *star reducible* if every element of W_c is star reducible to a product of commuting generators from S . Either X is a complete graph K_n with all labels $m(s, t) \geq 3$, or X appears in the list below, which comes from [6].






FIG. 2 The star reducible Coxeter groups

Definition 2.1.8 *If we have $m(s, t) \leq 3$ for all $s, t \in S$, we say W is simply laced.*

The following proposition is a subset of [7, Proposition 4.10]. Part (iii) of the original had a slight change and part (iv) was completely wrong. Note that a right-handed analogue of the proposition also holds.

Proposition 2.1.9 *Suppose that s and t are noncommuting generators of the Coxeter group W , and that $\tilde{t}_x \in \mathcal{L}_L^u$ whenever x is weakly complex, $ux \in W_c$ and $u \in S$. Then we have :*

(i)

$$\tilde{t}_s \tilde{t}_w \in \begin{cases} v\mathcal{L}_L^s & \text{if } sw < w, \\ \mathcal{L}_L^s & \text{if } sw > w; \end{cases}$$

(ii) $\tilde{t}_s \mathcal{L} \cap \mathcal{L} \subseteq \mathcal{L}_L^s$;

(iii) $\tilde{t}_s \mathcal{L}_L^t \subseteq \mathcal{L}_L^s$. ■

Chapter 3

Heaps of Pieces

3.1 Introduction

In Section 3.1, we introduce the basic properties of heaps. We will tend to follow Viennot's notation [13].

Definition 3.1.1 *Let P be a set equipped with a symmetric and reflexive binary relation $\mathcal{C}$. The elements of P are called pieces, and the relation $\mathcal{C}$ is called the concurrency relation. A labelled heap with pieces in P is a triple $(E, \leq, \varepsilon)$ where $(E, \leq)$ is a finite (possibly empty) partially ordered set with order relation denoted by $\leq$ and ε is a map $\varepsilon : E \rightarrow P$ satisfying the following two axioms:*

- (i) for every $a, b \in E$ such that $\varepsilon(a) \mathcal{C} \varepsilon(b)$, a and b are comparable in the order $\leq$;
- (ii) the order relation $\leq$ is the reflexive and transitive closure of the relation $\leq_{\mathcal{C}}$ such that for all $a, b \in E$, $a \leq_{\mathcal{C}} b$ if and only if both $a \leq b$ and $\varepsilon(a) \mathcal{C} \varepsilon(b)$.

Definition 3.1.2 *Let $(E, \leq, \varepsilon)$ and let $(E', \leq', \varepsilon')$ be two labelled heaps with pieces in P and the same concurrency relation, $\mathcal{C}$. An isomorphism $\phi : E \rightarrow E'$ of posets is said to be an isomorphism of labelled posets if $\varepsilon = \varepsilon' \circ \phi$.*

Definition 3.1.3 *A heap of pieces in P with concurrency relation $\mathcal{C}$ is an equivalence class of labelled heaps under labelled poset isomorphism. The set of such heaps is denoted $H(P, \mathcal{C})$. We denote the heap corresponding to the labelled heap $(E, \leq, \varepsilon)$ by $[E, \leq, \varepsilon]$. We will sometimes abuse notation by referring to a heap $[E, \leq, \varepsilon]$ simply as E .*

See Example 3.1.6 for an example of a labelled heap.

Definition 3.1.4 *The concurrency graph associated to the class of heaps $H(P, \mathcal{C})$ is the graph X whose vertices are the elements of P and for which there is an edge from $v \in P$ to $w \in P$ if and only if $v \neq w$ and $v \mathcal{C} w$. We will write $H(X)$ to mean $H(P, \mathcal{C})$.*

We are particularly interested in heaps arising from fully commutative elements in Coxeter groups as studied by Stembridge [12]. Elements of Coxeter groups give rise to heaps as follows.

Definition 3.1.5 *Let X be a graph, let S be the set of vertices of X and let $\mathcal{C}$ be the relation on S defined by $s_i \mathcal{C} s_j$ if and only if $s_i = s_j$ or s_i and s_j are adjacent vertices. Let $w = s_{i_1} s_{i_2} \cdots s_{i_l}$*

be an arbitrary word in the generators S of a Coxeter group $W = W(X)$. The word w gives a labelled heap $(E, \leq_E, \varepsilon)$ where $E = \{1, 2, \dots, l\}$, $\varepsilon(j) = s_{i_j}$, and the relation $\leq_{\mathcal{C}}$ of condition (ii) of Definition 3.1.1 can be defined by

$$a \leq_{\mathcal{C}} b \Leftrightarrow a \leq b \quad \text{and} \quad \varepsilon(a) \mathcal{C} \varepsilon(b),$$

where $\leq$ is the usual ordering on integers. The partial order $\leq_E$ is the reflexive and transitive closure of $\leq_{\mathcal{C}}$, and the heap of the word w is by definition the heap E_l corresponding to the given labelled heap.

The following example should give some insight into the previous definitions.

Example 3.1.6 Let X be the Coxeter graph of type $\tilde{C}_7$ as shown in Figure 3 below. Following the previous definition, $S = \{s_1, s_2, s_3, s_4, s_5, s_6, s_7, s_8\}$. Let $w = s_1 s_3 s_5 s_2 s_4 s_6 s_1 s_3 s_5 s_7 = s_{i_1} s_{i_2} \dots s_{i_9} s_{i_{10}}$. The word w gives a labelled heap E_{10} where $E = \{1, 2, \dots, 10\}$, and $\varepsilon(j) = s_{i_j}$. Note that s_{i_1} is the label for the vertex $s_1 \in X$ and s_{i_4} is the label for the vertex $s_2 \in X$. The vertices s_1 and s_2 are adjacent vertices in X so $s_{i_1} \leq_{\mathcal{C}} s_{i_4}$. The heap is shown in Figure 3(a).




FIG. 3. Coxeter graph of type $\tilde{C}_7$.




FIG 3(a). The heap of $w = s_1 s_3 s_5 s_2 s_4 s_6 s_1 s_3 s_5 s_7$ in type $\tilde{C}_7$.

Definition 3.1.7 Let $(E, \leq, \varepsilon)$ be a labelled heap with pieces in P and let F be a subset of E . Let ε' be the restriction of ε to F . Let $\mathcal{R}$ be the relation defined on F by $a \mathcal{R} b$ if and only if $a \leq b$ and $\varepsilon(a) \mathcal{C} \varepsilon(b)$. Let $\leq'$ be the transitive closure of $\mathcal{R}$. Then $(F, \leq', \varepsilon')$ is a labelled heap with pieces in P . The heap $[F, \leq', \varepsilon']$ is called a subheap of $[E, \leq, \varepsilon]$. If F is convex as a subset of E (if $x_i \leq x_j \leq x_k$ with $x_i, x_k \in F$, then $x_j \in F$), we call F a convex subheap of E . A trivial heap is a heap $[E, \leq, \varepsilon]$ for which the order relation $\leq$ is trivial, meaning that no element of E covers any other element.

Definition 3.1.8 Let $E = [E, \leq_E, \varepsilon]$ and $F = [F, \leq_F, \varepsilon']$ be two heaps in $H(P, \mathcal{C})$. We define the heap $G = [G, \leq_G, \varepsilon''] = E \odot F$ of $H(P, \mathcal{C})$ (which we call superposition of E over F as follows.

- (1) The underlying set G is the disjoint union of E and F .
- (2) The labelling map ε'' is the unique map $\varepsilon'' : G \rightarrow P$ whose restriction to E (respectively, F) is ε (respectively, ε').
- (3) The order relation $\leq_G$ is the transitive closure of the relation $\mathcal{R}$ on G , where $a \mathcal{R} b$ if and only

if one of the following three conditions holds:

- (i) $a, b \in E$ and $a \leq_E b$;
- (ii) $a, b \in F$ and $a \leq_F b$;
- (iii) $a \in E, b \in F$ and $\varepsilon(a) \mathcal{C} \varepsilon'(b)$.

3.2 The Boundary Map ∂

The following analogue to a certain boundary map ∂ in algebraic topology comes from [5].

Definition 3.2.1 Let k be a field. Let V_0 be the set of elements of $[E, \leq, \varepsilon]$, i.e., the set of elements of (a representative of) the underlying poset, E . We call the elements of V_0 vertices and denote their k -span by C_0 .

Let V_1 be the set of all pairs $(x, y) \in E \times E$ with $x < y$ and $\varepsilon(x) = \varepsilon(y)$ such that there is no element z for which we have both $\varepsilon(x) = \varepsilon(z) = \varepsilon(y)$ and $x < z < y$. We call the elements of V_1 edges and denote their k -span by C_1 .

For all other integers $i \in \mathbb{Z} \setminus \{0, 1\}$, we define $C_i = 0$.

The k -linear map $\partial = \partial_E : C_1 \rightarrow C_0$ is defined by its effect on the edges as follows:

$$\partial : (x, y) \mapsto \sum_{\substack{x < w < y \\ \varepsilon(w) \mathcal{C} \varepsilon(x)}} w.$$

Example 3.2.2 Consider the heap E arising from Example 3.1.6. The edges are $e_1 = (1, 7)$, $e_2 = (2, 8)$, and $e_3 = (3, 9)$. However, we will abuse notation and write $s_{i_k} = k$ so that the sums we create using the map ∂ make sense. Thus, $e_1 = (s_{i_1}, s_{i_7})$, $e_2 = (s_{i_2}, s_{i_8})$, and $e_3 = (s_{i_3}, s_{i_9})$. We now have $\partial(e_1) = s_{i_4}$, $\partial(e_2) = s_{i_4} + s_{i_5}$, and $\partial(e_3) = s_{i_5} + s_{i_6}$.

Definition 3.2.3 Let $E_n = E$ be a heap and let k be a field. If $v \in E$, we let $E(v)$ be the subheap of E obtained by defining $E(v) = E \setminus \{v\}$. We say E is acyclic if $\ker \partial_E = 0$. We say E is strongly acyclic if E is acyclic and $E(v)$ is acyclic for all $v \in E$. We say v is a boundary vertex of E if $v \in \text{Im}(\partial_E)$. We say v is an effective boundary vertex if $\partial_E(e_0) = v$ for an edge e_0 in E_n . Let R be the relation defined on the vertices of E_n by $v_1 R v_2$ if and only if $\partial_E(e_0) = v_1 + v_2$ for some edge e_0 . Linear equivalence is the reflexive, transitive closure of R . Since R is also symmetric, linear equivalence is an equivalence relation on the vertices of E . We will sometimes write R_E instead of R to show the relation holds on a particular set E .

Remark 3.2.4 The definitions for linear equivalence and effective boundary vertices are inspired by the notation and ideas of the Riemann–Roch theorem for graphs [1].

Example 3.2.5 In Example 3.1.6, since $\partial(e_1) = s_{i_4}$, we see that s_{i_4} is an effective boundary vertex. Furthermore, since $\partial(e_2) = s_{i_4} + s_{i_5}$ and $\partial(e_3) = s_{i_5} + s_{i_6}$, s_{i_4} and s_{i_5} are linearly equivalent and s_{i_5} and s_{i_6} are linearly equivalent. By the transitivity of linear equivalence, s_{i_6} and s_{i_4} are linearly equivalent. The equivalence classes are $\{s_{i_4}, s_{i_5}, s_{i_6}\}$, and the singletons $\{s_{i_1}\}$, $\{s_{i_2}\}$, $\{s_{i_3}\}$, $\{s_{i_7}\}$, $\{s_{i_8}\}$, $\{s_{i_9}\}$, $\{s_{i_{10}}\}$. It is clear that $\dim \text{Im}(\partial_E) = 3$. There are 3 edges, therefore $\ker \partial_E = 0$ and E is acyclic. If we consider the subheap $E(s_{i_5})$, we find that $\partial(e_1) = s_{i_4}$ and $\partial(e_2) = s_{i_4}$, thus $\ker \partial_{E(s_{i_5})} \neq 0$; hence the heap E is not strongly acyclic. Note that $\partial_E(e_2 - e_1) = (s_{i_4} + s_{i_5}) - (s_{i_4}) = s_{i_5}$, thus s_{i_5} is a boundary vertex; similarly, s_{i_6} is a boundary vertex.

3.3 Some Properties of Heaps and Full Commutativity

We recall property P1 from [5].

Definition 3.3.1 (Property P1) *Let $E = [E, \leq, \varepsilon]$ be a heap. We write $E(a) \prec^+ E$ (respectively, $E(a) \prec^- E$) if a is a maximal (respectively, minimal) vertex of E and there exists a maximal (respectively, minimal) vertex b of $E(a)$ with $\varepsilon(b) \neq \varepsilon(a)$ such that b is not maximal (respectively, minimal) in E . We write $E(a) \prec E$ if either $E(a) \prec^+ E$ or $E(a) \prec^- E$. If there is a (possibly trivial) sequence $E_1 \prec E_2 \prec \dots \prec E$ of heaps in $H(P, \mathbb{C})$ where E_1 is a trivial heap, we say that the heap E is dismantlable or that E has property P1.*

Example 3.3.2 *The heap arising from Example 3.1.6 is dismantlable and the following is a suitable chain. Note that $E(v_1, \dots, v_k)$ is the subheap corresponding to $E \setminus \{v_1, \dots, v_k\}$.*

$$\begin{aligned} E(s_{i_4}, s_{i_5}, s_{i_6}, s_{i_1}, s_{i_2}, s_{i_3}) &\prec E(s_{i_5}, s_{i_6}, s_{i_1}, s_{i_2}, s_{i_3}) \\ &\prec E(s_{i_6}, s_{i_1}, s_{i_2}, s_{i_3}) \\ &\prec E(s_{i_1}, s_{i_2}, s_{i_3}) \\ &\prec E(s_{i_2}, s_{i_3}) \\ &\prec E(s_{i_3}) \prec E \end{aligned}$$

where $E_1 = E(s_{i_4}, s_{i_5}, s_{i_6}, s_{i_1}, s_{i_2}, s_{i_3})$ is a trivial heap.

Let (W, S) be a Coxeter system with Coxeter graph X . Denote by S^* the free monoid on S . We call the elements of S *letters* and those of S^* *words*. Let $\phi_X : S^* \rightarrow W$ be the surjective morphism of monoid structures satisfying $\phi_X(i) = i$ for all $i \in S$. A word $i \in S^*$ is said to *represent* its image $w = \phi_X(i) \in W$; furthermore, if the length of i is minimal among the lengths of all the words that represent w , then i is a reduced expression for w . The *commutation monoid* $\text{Co}(X, S)$ is the quotient of the free monoid S^* by the congruence $\equiv$ generated by the commutation relations:

$$st \equiv ts \text{ for all } s, t \in S \text{ with } \phi_X(s)\phi_X(t) = \phi_X(t)\phi_X(s).$$

Note that, as a monoid, W is a quotient of $\text{Co}(X, S)$. We define $[i]$ to be the image of $i \in S^*$ in $\text{Co}(X, S)$. We will sometimes refer to the elements of $\text{Co}(X, S)$ as *traces*.

The product $\odot$ of heaps from Definition 3.1.8 is associative and $H(S, \mathbb{C})$ is a *monoid*, called the *heap monoid*, whose identity element is the empty heap.

We define the map $\phi_V : S^* \rightarrow H(S, \mathbb{C})$ by the relation

$$\text{for } w = \alpha_1 \alpha_2 \dots \alpha_n \in S^*, \phi_V(w) = \alpha_1 \odot \alpha_2 \odot \dots \odot \alpha_n \in H(S, \mathbb{C})$$

Proposition 3.3.3 *Let $H(S, \mathbb{C})$ be a heap monoid with pieces in S and concurrency relation $\mathbb{C}$. Let C be the complementary relation of $\mathbb{C}$. The morphism of monoids $\phi_V : S^* \rightarrow H(S, \mathbb{C})$ defined above induces an isomorphism $\bar{\phi}$ between the monoid $H(S, \mathbb{C})$ and the commutation monoid $\text{Co}(X, S)$.*

Proof. This is a restatement of [13, Proposition 3.1]. ■

Lemma 3.3.4 *Let $\text{Co}(X, S)$ be the commutation monoid defined above. Then the map $\psi : \text{Co}(X, S) \rightarrow W$ is a surjective morphism of monoid structures satisfying $\psi([s]) = \phi_X(s)$.*

Proof. The relations in $\text{Co}(X, S)$ are all relations in W , hence the map is well defined. $\blacksquare$

Theorem 3.3.5 *Maintain the above notation. The following diagram commutes, where all the maps are morphisms of monoids.*

$$\begin{array}{ccccc} S^* & \xrightarrow{\phi_V} & \text{Co}(X, S) & \xleftarrow{\bar{\phi}} & H(S, \mathcal{C}) \\ & \searrow \phi_X & \downarrow \psi & & \\ & & W & & \end{array}$$

Proof. The diagram collects the results from Proposition 3.3.3, Lemma 3.3.4, and the definitions preceding Proposition 3.3.3. $\blacksquare$

Definition 3.3.6 *The equivalence class $[\mathbf{s}] = \phi_V(\mathbf{s})$ of a given reduced word $\mathbf{s} \in S^*$ (relative to $\equiv$) consists of the words obtainable from $\mathbf{s}$ by transposing adjacent commuting pairs of generators from S . We call $[\mathbf{s}]$ the commutativity class of $\mathbf{s}$. If we consider the set $R(w) \subset S^*$ of all reduced expressions for some $w \in W$, $R(w)$ is a union of commutativity classes, namely the $\phi_V(R(w))$. If $\phi_V(R(w))$ is a singleton, we say that w is fully commutative. Note that this is a characterization of full commutativity. If $w \in W_c$, and $\mathbf{s}$ is a reduced expression for w , then $\phi_V(\mathbf{s})$ is independent of the choice of $\mathbf{s}$. We call $\bar{\phi}(\phi_V(\mathbf{s}))$ the heap of w .*

Definition 3.3.7 *Let E be a poset with partial order $\leq$. By a subposet of E , we mean a subset X of E equipped with the induced partial order; that is, the partial ordering of X such that for $x, y \in X$ we have $x \leq y$ in X if and only if $x \leq y$ in E . We define the closed interval $[x, y]_E$ to be $\{z \in E : x \leq z \leq y\}$. If X is totally ordered, then X is called a chain. A subposet X of E is convex if $y \in X$ whenever $x < y < z$ in E and $x, z \in X$. (In particular, an interval is convex.)*

Suppose that E is the heap of some $w \in W_c$. For each $s \in S$, the members of E with label s form a chain. We denote by $s^{(k)}$ the k -th least member of this chain with respect to E .

Some of the main results from [12] are summarized in the following theorem.

Theorem 3.3.8 *(i) The heap E of a word $\mathbf{s} \in S^*$ is the heap of some fully commutative element $w \in W$ if and only if the following two conditions hold:*

(a) there is no convex chain $i_1 < \dots < i_m$ in E such that $s_{i_1} = s_{i_3} = \dots = s$ and $s_{i_2} = s_{i_4} = \dots = t$, where $3 \leq m = m(s, t) < \infty$, and

(b) there is no covering relation $i < j$ in E such that $s_i = s_j$.

(ii) Subwords of fully commutative elements are fully commutative.

(iii) Let $s \in S$, and let $w \in W$ be fully commutative with heap E . If sw is not fully commutative, then sw is reduced and there is a unique $t \in S$ such that $m(s, t) \geq 3$ and $t^{(1)} < s^{(1)}$ in E . Moreover, $m(s, t) < \infty$ and

$$t^{(1)} < s^{(1)} < t^{(2)} < \dots < s^{(k)} \quad (\text{if } m(s, t) = 2k + 1)$$

$$t^{(1)} < s^{(1)} < t^{(2)} < \dots < t^{(k)} \quad (\text{if } m(s, t) = 2k),$$

is a convex chain in E .

Proof. For (i), see [12, Proposition 2.3]. For (ii), see [12, Prop 1.1]. For (iii), see [12, Lemma 3.1]. $\blacksquare$

Lemma 3.3.9 *Let E be a finite poset. Then $X \subset E$ is a convex subposet of E if and only if there is a sequence $E_0, E_1, \dots, E_k$ of subposets of E such that*

$$X = E_k \subset E_{k-1} \subset \dots \subset E_1 \subset E_0 = E$$

where $E_i \setminus E_{i+1} = \{\alpha\}$ with α maximal or minimal in E_i .

Proof. Assume that X is a convex subposet of E . The proof is by induction on k , where $k = |E \setminus X|$. If $k = 0$, then $X = E$ and there is nothing to prove. Suppose that $k > 0$. Since X is convex by assumption, X cannot contain all the maximal and minimal elements of E . Thus we can pick a maximal (or minimal) element $\alpha \in E$ such that $\alpha \notin X$. Since X is a convex subset of $E \setminus \{\alpha\}$ and $|(E \setminus \{\alpha\}) \setminus X| = k - 1$, we can apply the induction hypothesis. The conclusion follows.

Conversely, assume that there is a sequence of subposets of E such that

$$X = E_k \subset E_{k-1} \subset \dots \subset E_1 \subset E_0 = E$$

where $E_i \setminus E_{i+1} = \{\alpha\}$ with α maximal or minimal in E_i . Note that “is a convex subset of” is a transitive relation of subposets of a poset E . If we remove a maximal or minimal vertex $\alpha \in E$, we have a convex subset $E \setminus \{\alpha\}$. Combining these two facts gives us the result. ■

Lemma 3.3.10 *Let X be a convex subheap of a heap E . If e_0 is an edge in X , then e_0 is an edge in E .*

Proof. By way of contradiction, let $e_0 = (x, y)$ be an edge in X that is not an edge in E . Then there exists $z \in E$ such that $x < z < y$ and $\varepsilon(x) = \varepsilon(z) = \varepsilon(y)$, and z must be in X since X is a convex subset of E . Therefore e_0 is an edge in E . ■

Lemma 3.3.11 *Let E be a finite poset with convex subposet X . Then $[x, y]_X = [x, y]_E$.*

Proof. Let $z \in [x, y]_X$. Then $z \in X$ and $x \leq z \leq y$. Thus $z \in [x, y]_E$. Conversely, let $z \in [x, y]_E$ with $x \leq z \leq y$. Since $x, y \in X$, by convexity we have $z \in X$. Thus $z \in [x, y]_X$. ■

Lemma 3.3.12 *Let E be a heap and let X be a convex subheap of E . Let $e = (x, y)$ be an arbitrary edge in X . Identifying X as a subset of E , we have $\partial_E(e) = \partial_X(e)$.*

Proof. By definition of ∂_X , we have

$$\begin{aligned} \partial_X(e) &= \sum_{\substack{x < z < y \\ \varepsilon(z) \in \varepsilon(x) \\ z \in X}} z = \sum_{\substack{x < z < y \\ \varepsilon(z) \in \varepsilon(x) \\ z \in E}} z \\ &= \partial_E(e), \end{aligned}$$

where the equality of sums is by Lemma 3.3.11. ■

Lemma 3.3.13 *Let E be a heap and let X be a convex subheap of E .*

- (i) *If α, β are linearly equivalent vertices in X , then α, β are linearly equivalent in E .*
- (ii) *If α is an effective boundary vertex in X , then α is an effective boundary vertex in E .*

Proof. For (i), it is enough to assume that $\alpha R_X \beta$ in X . Then there is an edge of X , say e_0 , with $\partial_X(e_0) = \alpha + \beta$. By Lemma 3.3.10, e_0 is an edge of E . By Lemma 3.3.12, $\partial_E(e_0) = \alpha + \beta$. Therefore $\alpha R_E \beta$.

For (ii), let α be an effective boundary vertex in X . Then there is an edge e_0 in X with $\partial_X(e_0) = \alpha$. By Lemma 3.3.10, e_0 is an edge of E . By Lemma 3.3.12, $\partial_E(e_0) = \alpha$. Thus α is an effective boundary vertex of E . ■

Lemma 3.3.14 *Let W be a star reducible Coxeter group and let $w \in W_c$. Let E be the heap of w and let X be a convex subheap. Let B and B' be the sets of boundary vertices in E and X , respectively. Then $B' \subset B$.*

Proof. Assume for a contradiction that $\alpha \in B'$ and $\alpha \notin B$. By definition, there exist edges $e_0, e_1, \dots, e_r$ in X such that

$$\partial_X \left(\sum_{i=0}^r \lambda_i e_i \right) = \alpha.$$

By Lemma 3.3.12, $\partial_E(e_i) = \partial_X(e_i)$ for each $i = 0, 1, \dots, r$. Thus

$$\partial_X \left(\sum_{i=0}^r \lambda_i e_i \right) = \partial_E \left(\sum_{i=0}^r \lambda_i e_i \right) = \alpha. \quad \blacksquare$$

Lemma 3.3.15 *Let E be an arbitrary heap. If α, β are linearly equivalent in E and α is a boundary vertex, then β is a boundary vertex in E .*

Proof. We may assume that $\alpha R \beta$. By hypothesis, if $e_0, e_1, \dots, e_r$ are the edges of E , then $\partial_E(e_k) = \alpha + \beta$ for some k , and

$$\partial_E \left(\sum_{i=0}^r \lambda_i e_i \right) = \alpha$$

for some scalars $\lambda_1, \lambda_2, \dots, \lambda_r$. Hence

$$\beta = \partial(e_k) - \alpha = \partial(e_k) - \partial \left(\sum_{i=0}^r \lambda_i e_i \right) = \partial \left(e_k - \sum_{i=1}^r \lambda_i e_i \right),$$

which shows that β is a boundary vertex. ■

Definition 3.3.16 [5, Property P2] *We say a heap E has property P2 if it contains no convex chains of the form $x < y < z$ or $x < z$ with $\varepsilon(x) = \varepsilon(z)$ in either case.*

Example 3.3.17 *The heap arising from Example 3.1.6 does not have property P2. Although there are no convex chains of the form $s_{i_k} < s_{i_j}$ with $\varepsilon(s_{i_k}) = \varepsilon(s_{i_j})$, the chain $s_{i_1} < s_{i_4} < s_{i_7}$ violates the other requirement.*

The following theorem relates star reducible Coxeter groups, properties P1 and P2, and the definitions for strongly acyclic and acyclic heaps.

Theorem 3.3.18 *Let $W = W(X)$ be a star reducible Coxeter group with Coxeter graph X and suppose that E is the heap of (a reduced word for) some $w \in W_c$. Then*

- (i) *E has property P1 if and only if E is acyclic;*
- (ii) *if E is strongly acyclic, then E has property P2;*
- (iii) *if X is simply laced, then the converse to (ii) holds;*
- (iv) *any heap for $w \in W_c$ has property P1.*

Proof. Part (i) is [5, Theorem 2.4.4]. Part (ii) is [5, Proposition 2.2.7]. Part (iii) is [5, Theorem 2.4.2 (i)]. Part (iv) follows directly from the definition of star reducibility. ■

Lemma 3.3.19 *Let X be a bipartite, simply laced Coxeter graph with $X = X_1 \dot{\cup} X_2$, where X_1 and X_2 each consists of mutually nonadjacent vertices. Let $W = W(X)$ be the corresponding Coxeter group. Let E be the heap of some word on the Coxeter generators of W .*

- (i) *If $e = (x, y)$ and $\partial_E(e) = \sum_{j=0}^r \lambda_j \alpha_j$, then $x, y \in X_i$ for some $i \in \{1, 2\}$ and all α_j with $\lambda_j \neq 0$ lie in X_{3-i} .*
- (ii) *If E is acyclic and α is a boundary vertex of E , then $E \setminus \{\alpha\}$ is not acyclic.*
- (iii) *If E is the heap of a fully commutative element of W , then E has no boundary vertices.*

Proof. For (i), note that

$$\partial((x, y)) = \sum_{\substack{x < z < y \\ \varepsilon(x) \in \mathcal{C} \varepsilon(z)}} z.$$

Since (x, y) is an edge, $\varepsilon(z)$ is adjacent to $\varepsilon(x)$. Therefore, if $\varepsilon(x) = \varepsilon(y) \in X_i$, then $\varepsilon(z) \in X_{3-i}$. The assertion follows.

For (ii), assume that α is a boundary vertex of E . By definition, there exist edges $e_0, e_1, \dots, e_r$ in E such that

$$\partial_E \left(\sum_{i=0}^r \lambda_i e_i \right) = \alpha.$$

By (i), α is not an endpoint of any edge e_i , so deleting α gives us

$$\partial_{E \setminus \{\alpha\}} \left(\sum_{i=0}^r \lambda_i e_i \right) = 0.$$

But then $\ker \partial_{E \setminus \{\alpha\}} \neq 0$ and hence $E \setminus \{\alpha\}$ is not acyclic.

For (iii), assume by way of contradiction that α is a boundary vertex in E . By definition, there exist edges $e_0, e_1, \dots, e_r$ in E such that

$$\partial_E \left(\sum_{i=0}^r \lambda_i e_i \right) = \alpha.$$

By (i), α is not an endpoint of any edge e_i . Since X is simply laced, E has Property P2 and by Theorem 3.3.18 (iii), E is strongly acyclic. By (ii), if we delete the vertex α , we have that $E \setminus \{\alpha\}$ is not acyclic and hence E is not strongly acyclic, a contradiction. ■

3.4 Main Theorem

Theorem 3.4.1 *Let X be a bipartite Coxeter graph with star reducible Coxeter group $W = W(X)$. In the heap E of a fully commutative element $w \in W$, every boundary vertex is linearly equivalent to an effective boundary vertex.*

Proof. If X is simply laced, then the claim is true for $W(X)$ by Lemma 3.3.19 (iii). By the classification of star reducible Coxeter groups in [6], we can assume X is a straight line graph. Let $x \in W_c$, and let E be the heap of x as in Definition 3.3.6.

The proof is by induction on n , the number of vertices in the heap E . If x is a product of commuting generators from S , then the heap of x has no edges and hence no boundary vertices. Therefore the claim is true vacuously and the cases $n = 0$ and $n = 1$ are covered.

Since W is a star reducible Coxeter group, we may assume that x either has a reduced expression beginning with st or ending with ts , where $I = \{s, t\}$ is a pair of noncommuting generators in S . We deal here with the case where x has a reduced expression beginning with st . If x has a reduced expression ending in ts , a symmetrical argument gives the same conclusion.

Taking star operations with respect to I , we can star reduce $x = stv$ (reduced) to $w = tv$. Since w is a subword of the fully commutative element x , by Theorem 3.3.8 (ii), w is fully commutative. Since the heap E_{n-1} of w is a convex subheap of E_n , by Lemma 3.3.14, the boundary vertices B' of E_{n-1} are a subset of the boundary vertices B of E_n . If $B' = B$, by Lemma 3.3.13 the claim is true.

Suppose there is a boundary vertex $\alpha \in B$ with $\alpha \notin B'$. Then there exist edges $e_0, e_1, \dots, e_r$ in E_n such that

$$\partial_E \left(\sum_{i=0}^r \lambda_i e_i \right) = \alpha.$$

where $e_0 = (s^{(0)}, s^{(1)})$ with $s^{(0)} = s$ in the definition of x . Because $t^{(1)}$ is minimal in $E \setminus \{s^{(0)}\}$, $t^{(1)}$ does not appear in $\partial_E(e_i)$ for any $i \neq 0$. Thus $t^{(1)} = \alpha$, $\lambda_0 = 1$, and $t^{(1)}$ is unique. We want to show that $t^{(1)}$ is linearly equivalent to a boundary vertex $\beta \in B'$.

Since $t^{(1)}$ is a boundary vertex in E_n , $E_n(t^{(1)})$ has a cycle by Lemma 3.3.19 (ii). By Theorem 3.3.18 (i), $E_n(t^{(1)})$ does not have Property P1. Therefore $E_n(t^{(1)})$ is not the heap of a fully commutative element by Theorem 3.3.18 (iv). By Theorem 3.3.8 (i), either $E_n(t^{(1)})$ has a convex chain $s^{(0)} < u^{(1)} < s^{(1)} < \dots < u^{(k)}$ (or $s^{(k)}$, depending on $m(s, u)$) or a convex chain $s^{(0)} < s^{(1)}$. If the second statement is true, then $s^{(0)} < t^{(1)} < s^{(1)}$ in E_n , $t^{(1)}$ is an effective vertex in E_n and we are done.

If the first statement is true, then we must have $\partial_E(s^{(0)}, s^{(1)}) = u^{(1)} + t^{(1)}$ and $u^{(1)}$ is a boundary vertex in E_n , by Lemma 3.3.15. As $t^{(1)}$ is the only boundary vertex in B that is not in B' , $u^{(1)}$ must be a boundary vertex in B' . By induction, $u^{(1)}$ is linearly equivalent in E_{n-1} to an effective boundary vertex in E_{n-1} and thus in E_n by Lemma 3.3.13. As $t^{(1)}$ is linearly equivalent to $u^{(1)}$ in E_n , by the transitivity of linear equivalence, $t^{(1)}$ is linearly equivalent in E_n to an effective boundary vertex in E_n , and the claim is true. ■

The following is an example to illustrate that the theorem holds.

Example 3.4.2 *Let X be the Coxeter graph of type $\tilde{C}_7$, as shown in Figure 5 below. Since X has no odd cycles, we see that X is bipartite. Also note that $W(X)$ is star reducible by the classification in [6]. Let $w = s_1 s_3 s_5 s_2 s_4 s_6 s_1 s_3 s_5 s_7 = s_{i_1} s_{i_2} \dots s_{i_9} s_{i_{10}}$ and let E be the heap of w . In Example*

3.2.5, we showed that the boundary vertices of E are $s_{i_4}, s_{i_5}, s_{i_6}$. Because $\partial(e_1) = s_{i_4}$, we see that s_{i_4} is an effective boundary vertex. Furthermore, since $\partial(e_2) = s_{i_4} + s_{i_5}$ and $\partial(e_3) = s_{i_5} + s_{i_6}$, s_{i_4} and s_{i_5} are linearly equivalent and s_{i_5} and s_{i_6} are linearly equivalent. By the transitivity of linear equivalence, s_{i_6} and s_{i_4} are linearly equivalent. Thus s_{i_5} and s_{i_6} are both linearly equivalent to the effective boundary vertex s_{i_4} and the theorem holds.



FIG. 5. Coxeter graph of type $\tilde{C}_7$.




FIG 5(a). The heap of $w = s_1 s_3 s_5 s_2 s_4 s_6 s_1 s_3 s_5 s_7$ in type $\tilde{C}_7$.

The following is a non-example.

Example 3.4.3 Consider the bipartite Coxeter graph of Figure 6.




FIG. 6. Coxeter graph on six vertices.

Let $S = \{s_1, s_2, \dots, s_6\}$. Let $w = s_1 s_3 s_5 s_2 s_4 s_1 s_3 s_2 s_4 s_6 s_1 s_3 s_5 = s_{i_1} s_{i_2} \dots s_{i_{12}} s_{i_{13}}$. The word w gives a labelled heap E_{13} where $E = \{1, 2, \dots, 13\}$, $\varepsilon(j) = s_{i_j}$. The heap is shown in Figure 7.




FIG 7. The labelled heap of $w = s_1 s_3 s_5 s_2 s_4 s_1 s_3 s_2 s_4 s_6 s_1 s_3 s_5$.

Note that w is fully commutative and the heap is dismantlable. However $s_{i_{10}}$ is a boundary vertex that is not linearly equivalent to an effective boundary vertex. Theorem 3.4.1 does not apply in this case because the group defined by the Coxeter graph is not one of the star reducible Coxeter groups.

Chapter 4

Property W

4.1 Introduction

In this chapter we present an application of Theorem 3.4.1.

Lemma 4.1.1 *Let E be a heap and let $\alpha \in E$. Then $|\dim \ker \partial_E - \dim \ker \partial_{E(\alpha)}| \leq 1$*

Proof. This is [5, Theorem 2.1.1].

The reader may wish to review the definitions of generalized Temperley–Lieb algebra and $\tilde{t}$ -basis given in §2.1.

Definition 4.1.2 *Let W be a Coxeter group with Coxeter graph X and let $w \in W_c$. Let $w = s_1 s_2 \cdots s_r$ be a reduced expression for w . For each $s \in S$, let $b_s = v^{-1} \tilde{t}_1 + \tilde{t}_s$, then define $b_w \in TL(X)$ by $b_w = b_{s_1} b_{s_2} \cdots b_{s_r}$. Note that the element b_w is well-defined because any two reduced expressions for w are commutation equivalent and $b_s b_t = b_t b_s$ if $st = ts$. The set $\{b_w : w \in W_c\}$ is a basis for $TL(X)$, called the monomial basis; see Lemma 4.1.5 (ii) for more details.*

Definition 4.1.3 *Let $W = W(X)$ be a star reducible Coxeter group. We define the function $h : S^* \rightarrow \mathbb{Z}^{\geq 0}$ by $h(\mathbf{u}) = \dim \ker \partial_{\phi(\phi_V(\mathbf{u}))}$. We sometimes write $h([\mathbf{u}])$ instead of $h(\mathbf{u})$; this makes sense by Theorem 3.3.5.*

Lemma 4.1.4 *Let $W(X)$ be a star reducible Coxeter group, let $\mathbf{s} = s_1 s_2 \cdots s_r \in S^*$, and let $b(\mathbf{s}) \in TL(X)$ given by $b(\mathbf{s}) = b_{s_1} b_{s_2} \cdots b_{s_r}$. Express b as a linear combination of the monomial basis, namely*

$$b(\mathbf{s}) = \sum_{w \in W_c} \lambda_w b_w.$$

Then each λ_w is an integer multiple of $(v + v^{-1})^{h(\mathbf{s})}$.

Proof. This is [6, Lemma 2.9]. ■

Recall from §1 that $\mathcal{A}^-$ is the ring $\mathbb{Z}[v^{-1}]$.

Lemma 4.1.5 *Let $W = W(X)$ be a star reducible Coxeter group with Coxeter graph X .*

- (i) *If $x \in W$ is weakly complex, then $\tilde{t}_x \in \mathcal{L}$.*
- (ii) *The b -basis and the c -basis of $TL(X)$ have the same $\mathbb{Z}$ -span. In particular, the b -basis is an $\mathcal{A}^-$ -basis for $\mathcal{L}$.*

Proof. Part (i) is [6, Lemma 3.8(i)]. Part (ii) is [6, Lemma 2.10]. ■

Lemma 4.1.6 *Let $\mathbf{s} = s_1 s_2 \cdots s_r \in S^*$ with $\overline{\phi}(\phi_V(\mathbf{s}))$ acyclic and write*

$$\tilde{t}_{\mathbf{s}} = \tilde{t}_{s_1} \tilde{t}_{s_2} \cdots \tilde{t}_{s_r}.$$

Let $\tilde{t}_{\mathbf{s}'}$ be the result of replacing k distinct $\tilde{t}_{s_i}$ with $(-v^{-1})$. Then $\tilde{t}_{\mathbf{s}'} \in \mathcal{L}$.

Proof. Let $\mathbf{s} = s_1 s_2 \cdots s_r$ be a reduced expression for $w \in W_c$ and write

$$\begin{aligned} \tilde{t}_{\mathbf{s}} &= \tilde{t}_{s_1} \tilde{t}_{s_2} \cdots \tilde{t}_{s_r} \\ &= (b_{s_1} - v^{-1})(b_{s_2} - v^{-1}) \cdots (b_{s_r} - v^{-1}). \end{aligned}$$

Expanding the parentheses, we can express $\tilde{t}_w$ as a linear combination of 2^r elements $(-v)^{-k}b(\mathbf{u})$, where $\mathbf{u}$ is obtained from $\mathbf{s}$ by deletion of k generators. We have

$$(-v)^{-k}b(\mathbf{u}) = \sum_{w \in W_c} (-v)^{-k} \lambda_w b_w.$$

By Lemma 4.1.4, $\lambda_w \in v^{h(\mathbf{u})}\mathcal{A}^-$. By Lemma 4.1.1, $h(\mathbf{u}) \leq h(\mathbf{s}) + k$. Since $\overline{\phi}(\phi_V(\mathbf{s}))$ is acyclic by hypothesis, $h(\mathbf{s}) = 0$. Thus $h(\mathbf{u}) \leq k$ and $(-v^{-k})\lambda_w \in v^{h(\mathbf{u})-k}\mathcal{A}^- \subset \mathcal{A}^-$. By Lemma 4.1.5 (ii), $b_w \in \mathcal{L}$ so $(-v^{-k})\lambda_w b_w \in \mathcal{L}$ and $(-v^{-k})b(\mathbf{u}) \in \mathcal{L}$.

Now, $\tilde{t}_{\mathbf{s}'}$ is a linear combination of 2^{r-k} terms of the form $(-v^{-k})b(\mathbf{u})$, and the conclusion follows. ■

Lemma 4.1.7 *Let W be a Coxeter group and let $I = \{s, t\}$ be a pair of noncommuting generators in S . Suppose that whenever x is weakly complex, $ux \in W_c$ and $u \in S$, we have $t_x \in \mathcal{L}_L^u$. Let $w = w_I w^I$ be such that $w^I \in W_c$.*

(i) If $sw_I < w_I$, then $\tilde{t}_w \in \mathcal{L}_L^s$.

(ii) If $w_I = w_{st}$, the longest element in W_I , then $\tilde{t}_{w_I} \tilde{t}_{w^I} \in v^{-1}\mathcal{L}_L^s$.

Proof. This is [7, Lemma 4.11]. A right-handed analogue of this lemma also holds. ■

4.2 Acyclic Case

Lemma 4.2.1 *Let $W(X)$ be a star reducible Coxeter group with a straight line Coxeter graph X and let $\mathbf{s}$ be a reduced expression for $w \in W_c$. If $x = sw \notin W_c$ and the heap $E = \overline{\phi}(s[\mathbf{s}])$ is acyclic, then $\tilde{t}_x \in v^{-1}\mathcal{L}$.*

Proof. Since X is a straight line, each Coxeter generator fails to commute with at most two other generators. We will use this fact freely. The proof is by induction on $\ell(w)$. There are no weakly complex elements of length 0, thus the base case is vacuous. By Theorem 3.3.8 (iii) it follows that sw is reduced. From the proof of Theorem 3.3.8 (i), there exists a unique $t \in S$ such that $3 \leq m(s, t) < \infty$ and we have

$$s^{(1)} < t^{(1)} < s^{(2)} < t^{(2)} < \cdots < s^{(k+1)} \quad (\text{if } m(s, t) = 2k + 1)$$

$$s^{(1)} < t^{(1)} < s^{(2)} < t^{(2)} < \cdots < s^{(k)} < t^{(k)} \quad (\text{if } m(s, t) = 2k),$$

occurring as a convex chain c in the heap $\overline{\phi}(s[s])$, where s is a reduced expression for w .

We first show that the five cases that follow are exhaustive. Since E is acyclic, by Theorem 3.3.18 (i), E has property P1. As sw is weakly complex, E is not trivial, so E can be left or right star reduced. Thus there exist $\alpha, \beta \in E$ such that $E(\alpha) \prec^+ E$ (respectively, $E(\alpha) \prec^- E$) if α is a maximal (respectively, minimal) vertex of E and there exists a maximal (respectively, minimal) vertex β of $E(\alpha)$ with $\varepsilon(\beta) \neq \varepsilon(\alpha)$ such that β is not maximal (respectively, minimal) in E . This gives us two subcases for possible star reductions involving α and β .

If $E(\alpha) \prec^- E$, then there exist elements $a, b \in S$ such that $a = \varepsilon(\alpha)$ and $b = \varepsilon(\beta)$. Either $\alpha \notin \mathbf{c}$ or $\alpha \in \mathbf{c}$. If $\alpha \notin \mathbf{c}$, then $\beta \notin \mathbf{c}$ because β is minimal in $E(\alpha)$. This is case (i) below. If $\alpha \in \mathbf{c}$, then $\alpha = s^{(1)}$. By the convexity of $\mathbf{c}$, $\beta = t^{(1)}$ and $\beta \in \mathbf{c}$; this is case (ii) below.

Suppose on the other hand that $E(\alpha) \prec^+ E$. Either $\alpha, \beta \in \mathbf{c}$, $\alpha, \beta \notin \mathbf{c}$, or $\alpha \notin \mathbf{c}$ but $\beta \in \mathbf{c}$. These are dealt with respectively in cases (iii), (iv), and (v) below.

Case (i) In this case, both a and b commute with each of s and t and we have $w = abw'$ reduced. Since x is weakly complex, by Lemma 4.1.5 (i), $\tilde{t}_x \in \mathcal{L}$ and $\ell(sw') > \ell(w')$. Also, since $x = sw = absw'$ is reduced, we can left star reduce x to the element bsw' . Now, bsw' and sw' are both weakly complex since α and β are not involved in the chain c . Thus the induction hypothesis is satisfied and we have $\tilde{t}_{bsw'} \in v^{-1}\mathcal{L}$ and $\tilde{t}_{sw'} \in v^{-1}\mathcal{L}$. Therefore $v\tilde{t}_{sw'} \in \mathcal{L}$ and

$$\tilde{t}_b(v\tilde{t}_{sw'}) = v\tilde{t}_{bsw'} \in \mathcal{L}^b$$

by Lemma 4.1.5 (i) and Proposition 2.1.9 (ii). By Proposition 2.1.9 (iii), $\tilde{t}_a(v\tilde{t}_{bsw'}) \in \mathcal{L}^a$. Hence

$$\tilde{t}_a\tilde{t}_{bsw'} = \tilde{t}_{absw'} = \tilde{t}_x \in v^{-1}\mathcal{L}^a \subset v^{-1}\mathcal{L}.$$

Case (ii) In this case, $a = s$, $b = t$, and we have $x = abw'$ reduced. We have $s[s] = [\mathbf{c}\mathbf{u}] \in \text{Co}(X, S)$ with $\alpha, \beta \in \mathbf{c}$. Thus $sw = w_{st}w'$ reduced, where w_{st} is the longest element in the parabolic subgroup generated by $I = \{s, t\} = \{a, b\}$ and $\mathcal{L}(w') \cap I = \emptyset$. Therefore

$$\tilde{t}_{sw} = \tilde{t}_{w_{st}w'} = \tilde{t}_{w_{st}}\tilde{t}_{w'}$$

The hypothesis of Lemma 4.1.7 (ii) is satisfied and we have

$$\tilde{t}_{w_{st}}\tilde{t}_{w'} \in v^{-1}\mathcal{L}^s \subset v^{-1}\mathcal{L}$$

and the claim is true.

Case (iii) In this case, $\{s, t\} = \{a, b\}$ and $\alpha, \beta \in \mathbf{c}$ are such that $s[s] = [\mathbf{u}\mathbf{c}] \in \text{Co}(X, S)$. We can now argue as in case (ii) and the result follows.

Case (iv) In this case, $x = sw = sw'ba$ reduced and neither α nor β is in c . We can right star reduce x to the element $sw'b$. Now, $sw'b$ and sw' are both weakly complex since $\alpha, \beta \notin \mathbf{c}$. Thus the induction hypothesis is satisfied and we have $\tilde{t}_{sw'b} \in v^{-1}\mathcal{L}$ and $\tilde{t}_{sw'} \in v^{-1}\mathcal{L}$. Therefore $v\tilde{t}_{sw'} \in \mathcal{L}$ and

$$(v\tilde{t}_{sw'})\tilde{t}_b = v\tilde{t}_{sw'b} \in \mathcal{L}^b$$

by Lemma 4.1.5 (i) and Proposition 2.1.9 (ii). By Proposition 2.1.9 (iii), $(v\tilde{t}_{sw'b})\tilde{t}_a \in \mathcal{L}^a$. Hence

$$\tilde{t}_{sw'b}\tilde{t}_a = \tilde{t}_{sw'ba} = \tilde{t}_x \in v^{-1}\mathcal{L}^a \subset v^{-1}\mathcal{L}$$

and the claim is true.

Case (v) In this case, $\{s, t\} = \{b', b\}$ for $b \neq a$, and $x = sw = u'w_{st}u''a = x'ba$ reduced. We have $\beta \in c$ but $\alpha \notin c$ and $s[s] = [u\alpha] \in \text{Co}(X, S)$. Since w_{st} is the longest element in the parabolic subgroup $\langle s, t \rangle$, we can use the relation

$$\tilde{t}_{w_{st}} = - \sum_{\substack{u \in \langle s, t \rangle \\ u < w_{st}}} v^{\ell(u) - m(s, t)} \tilde{t}_u$$

to write

$$\begin{aligned} \tilde{t}_x &= \tilde{t}_{u'} \tilde{t}_{w_{st}} \tilde{t}_{u''} \tilde{t}_a \\ &= -\tilde{t}_{u'} \left(v^{-1} \tilde{t}_{w_{st}} + v^{-1} \sum_{u \leq sw_{st}} v^{\ell(u) - \ell(sw_{st})} \tilde{t}_u \right) \tilde{t}_{u''} \tilde{t}_a. \end{aligned}$$

By Lemma 4.1.6, we have $\tilde{t}_u \tilde{t}_u \tilde{t}_{u''} \tilde{t}_a \in \mathcal{L}$ where u is as in the sum above. To see that $\tilde{t}_{u'} \tilde{t}_{w_{st}} \tilde{t}_{u''} \tilde{t}_a \in \mathcal{L}$, first note that $u'tw_{st}u''$ is an expression for x' , which is fully commutative. If x' has a reduced expression ending in a , since $x'b$ is weakly complex, by Theorem 3.3.8 (i) the only element in $\mathcal{R}(x')$ not commuting with b is b' . But then $a = b'$ and $[ca]$ is a convex chain of length $m(s, t) + 1$ and $s[s]$ is not reduced, a contradiction. Therefore x' has no reduced expression ending in a , $x'a$ is either fully commutative or weakly complex, and by Lemma 4.1.5 (i), $\tilde{t}_{x'a} \in \mathcal{L}$. Putting all this together, we have $\tilde{t}_x \in v^{-1}\mathcal{L}$. $\blacksquare$

4.3 Forbidden Configurations

The following is a collection of combinatorial results concerning inadmissible configurations in heaps of fully commutative elements. For all of §4.3, let X be a straight line graph with vertices $s_1, s_2, \dots, s_n$ such that s_i, s_j are adjacent if and only if $|i - j| = 1$.

Lemma 4.3.1 *Let X be a Coxeter graph of type B_3 and assume $m(s_1, s_2) = 4$. If $w \in W_c(X)$, then a reduced expression for w has at most three occurrences of each of s_2 and s_3 .*

Proof. It follows from [9, §1.8] that we have $w < w_0$ in the Bruhat order. This means that given a reduced expression $\mathbf{s}$ for w_0 , some subexpression of w_0 is equal to w , by [9, §5.10]. One such reduced expression for w_0 is $\mathbf{s} = s_1 s_3 s_2 s_1 s_3 s_2 s_1 s_3 s_2$. Since every expression for $w \in W_c$ has the same number of generators of each type, and $\mathbf{s}$ contains three occurrences of s_2 and three occurrences of s_3 , the result follows. $\blacksquare$

Recall from §3.3 and Theorem 3.3.5 that each heap E is associated to a unique trace $\overline{\phi}^{-1}(E)$, which we call the trace of E .

Lemma 4.3.2 *Let X be a straight line Coxeter graph and let E be the heap of $w \in W_c(X)$.*

- (i) *If $m(s_j, s_{j+1}) = 3$ for all $j \geq i$ and $[x s_i y s_i z]$ is a trace for E , then s_{i-1} must occur in $\mathbf{y}$.*
- (ii) *If $m(s_j, s_{j+1}) = 3$ for all $j \leq i$ and $[x s_i y s_i z]$ is a trace for E , then s_{i+1} must occur in $\mathbf{y}$.*

Proof. Without loss of generality, we may assume that there are no occurrences of s_i in $\mathbf{y}$. We prove (i) by induction on $n - i$; (ii) follows similarly. If $i = n$, then we have $\overline{\phi}^{-1}(E) = [x s_n y s_n z]$. Since there are no generators s_j such that $j > n$, if there is no occurrence of s_{i-1} in $\mathbf{y}$, $x s_n y s_n z$ is not reduced, which completes the base case.

Assume that $i < n$. Assume by way of contradiction that there are no occurrences of s_{i-1} in $\mathbf{y}$. Since the only other generator that does not commute with s_i is s_{i+1} , there must be at least one occurrence of s_{i+1} in $\mathbf{y}$, else $\mathbf{x}s_i\mathbf{y}s_i\mathbf{z}$ would not be reduced. If there is only one occurrence of s_{i+1} in $\mathbf{y}$, we have a contradiction to Theorem 3.3.8 (i) because $m(s_i, s_{i+1}) = 3$. If there is more than one occurrence of s_{i+1} in $\mathbf{y}$, then we are done by induction because there exist two occurrences of s_{i+1} with no s_i between them. ■

Lemma 4.3.3 *Let X be a Coxeter graph of type F_n and let E be the heap of $w \in W_c(X)$. Assume that $m(s_2, s_3) = 4$.*

(i) *If $[\mathbf{x}s_3\mathbf{y}s_3\mathbf{z}]$ is the trace of E and s_3, s_4 are not in $\mathbf{y}$, then $\mathbf{y}$ contains a unique s_2 .*

The heap E has no trace of the following forms:

(ii) $[\mathbf{x}(s_1s_2)\mathbf{y}(s_3s_2s_1s_3)\mathbf{z}]$ in which $s_3, s_4 \notin \mathbf{y}$;

(iii) $[\mathbf{x}(s_4s_3)\mathbf{y}(s_2s_3s_2s_4)\mathbf{z}]$ in which $s_4, s_5 \notin \mathbf{y}$;

(iv) $[\mathbf{x}(s_3s_1s_2s_3)\mathbf{y}(s_2s_1)\mathbf{z}]$ in which $s_3, s_4 \notin \mathbf{y}$;

(v) $[\mathbf{x}(s_4s_2s_3s_2)\mathbf{y}(s_3s_4)\mathbf{z}]$ in which $s_4, s_5 \notin \mathbf{y}$.

Proof. For part (i), $\mathbf{y}$ must contain an occurrence of s_2 for the expression to be reduced. If $\mathbf{y}$ contains two occurrences of s_2 , this contradicts Lemma 4.3.2 (ii).

For part (ii), we may assume $\mathbf{y}$ is as short as possible, which means that only s_1 and s_2 can appear in $\mathbf{y}$, since by assumption, $\mathbf{y}$ has no occurrences of s_3 or s_4 . If $\mathbf{y}$ is empty, we have a subword of the form $s_2s_3s_2s_1s_3$ which contradicts Theorem 3.3.8 (i). If the rightmost generator of $\mathbf{y}$ is s_1 , we have $s_1s_3s_2s_1s_3$, which also contradicts Theorem 3.3.8 (i). If the rightmost generator of $\mathbf{y}$ is s_2 , we have $s_2s_3s_2s_1s_3$, which contradicts Theorem 3.3.8 (i) as well.

For part (iii), we may assume $\mathbf{y}$ is as short as possible, which means that only s_1, s_2 and s_3 can appear in $\mathbf{y}$, since by assumption, $\mathbf{y}$ has no occurrences of s_4 or s_5 . If there is an occurrence of s_3 in $\mathbf{y}$, then by part (i), $\mathbf{y}$ has no occurrences of s_2 to the right of the rightmost occurrence of s_3 . But now the subword $s_3s_2s_3s_2$ contradicts Theorem 3.3.8 (i), since $m(s_2, s_3) = 4$, thus there is no occurrence of s_3 in $\mathbf{y}$. By part (i) again, there is no occurrence of s_2 , either. But then $\mathbf{y} = s_1$, yielding another contradiction to Theorem 3.3.8 (i).

Parts (iv) and (v) are handled by symmetrical arguments to parts (ii) and (iii), respectively. The conclusion follows. ■

Lemma 4.3.4 *Let X be a Coxeter graph of type F_n and let E be the heap of $w \in W_c(X)$. Assume that $m(s_2, s_3) = 4$. Then E cannot have a trace of the following form*

$$[\mathbf{x}(s_i s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2} s_i) \mathbf{z}]$$

in which $\mathbf{y}$ has no occurrences of s_{i+2} or s_{i+3} .

Proof. The proof is by induction on i . Assume that $i = 1$, and that E does have such a trace. Then

$$[\mathbf{x}(s_1 s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2} s_i) \mathbf{z}] = [\mathbf{x}(s_3 s_1 s_2) \mathbf{y}(s_2 s_1 s_3) \mathbf{z}]$$

where $s_3, s_4 \notin \mathbf{y}$. We may assume $\mathbf{y}$ is as short as possible, which means that only s_1 and s_2 can appear in $\mathbf{y}$. Now, $\mathbf{y}$ is non-empty, else $[\mathbf{x}(s_3 s_1 s_2) \mathbf{y}(s_2 s_1 s_3) \mathbf{z}]$ would not be reduced. Since $s_1 s_2 \mathbf{y} s_2 s_1$ has length greater than 4 and $m(s_1, s_2) = 3$, $s_1 s_2 \mathbf{y} s_2 s_1$ is not reduced, a contradiction.

Assume that $i = 2$, and that E does have such a trace. Then

$$[\mathbf{x}(s_i s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2} s_i) \mathbf{z}] = [\mathbf{x}(s_4 s_2 s_3) \mathbf{y}(s_3 s_2 s_4) \mathbf{z}]$$

where $s_4, s_5 \notin \mathbf{y}$. We may assume that $\mathbf{y}$ is as short as possible, which means that only s_1, s_2 and s_3 can appear in $\mathbf{y}$. By Lemma 4.3.1, $s_2 s_3 \mathbf{y} s_3 s_2$ has at most three occurrences of each of s_2 and s_3 . Suppose for a contradiction that there is an occurrence of s_3 in $\mathbf{y}$. In this case, since there are no occurrences of s_4 or s_5 by hypothesis, there must be at least two occurrences of s_2 in $\mathbf{y}$. However, since there are now four occurrences of s_2 in $s_2 s_3 \mathbf{y} s_3 s_2$, this contradicts Lemma 4.3.1. Thus it must be the case that there are no occurrences of s_3 in $\mathbf{y}$. By Lemma 4.3.3 (i), $\mathbf{y}$ contains a unique s_2 . But then there must be two occurrences of s_1 in $\mathbf{y}$, else the alternating product of s_2 and s_3 corresponds to a convex chain. In fact, we must have $\mathbf{y} = s_1 s_2 s_1$. But then $s_2 s_3 s_1 s_2 s_1 s_3 s_2$ is not fully commutative since $m(s_1, s_2) = 3$, a contradiction.

Assume now that $i > 2$, and that E does have such a trace. Note also that $m(s_i, s_{i+1}) = 3$. Then E has a trace of the form $[x s_i s_{i+2} s_{i+1} \mathbf{y}' s_{i+1} \mathbf{z}']$ in which $\mathbf{y}'$ has no occurrences of s_{i+1} . Since $\mathbf{y}'$ is a subword of $\mathbf{y}$, $\mathbf{y}'$ cannot contain any occurrences of s_{i+2} and s_{i+3} . Since $\mathbf{y}'$ contains no occurrences of s_{i+1} , it must be the case that $\mathbf{y}'$ contains at least two occurrences of s_i , or we would contradict Theorem 3.3.8 (i). There must also be an occurrence of s_i in $\mathbf{z}'$, because of the subword $[s_{i+1} s_{i+2} s_i \mathbf{z}]$ in the original form of the trace. We can now write the trace as

$$[x s_i s_{i+2} s_{i+1} \mathbf{y}_1 s_i \mathbf{y}_2 s_i \mathbf{y}_3 s_{i+1} \mathbf{z}_1 s_i \mathbf{z}_2],$$

where $\mathbf{y}_1, \mathbf{y}_3$ and $\mathbf{z}_1$ contain no occurrences of s_i , and $\mathbf{y}_1, \mathbf{y}_2$ and $\mathbf{y}_3$ contain no occurrences of s_{i+1}, s_{i+2} or s_{i+3} . Furthermore, s_{i+1} cannot appear in $\mathbf{z}_1$, or else we would have $[s_{i+1} \mathbf{z}_1 s_i] = [s_{i+1} \mathbf{z}'_1 s_{i+1} \mathbf{z}']$ which contradicts Lemma 4.3.2 (i) applied to $[s_{i+1} \mathbf{z}'_1 s_{i+1}]$.

Applying the relation $[\mathbf{y}_3 s_{i+1}] = [s_{i+1} \mathbf{y}_3]$, the trace is equal to

$$[x s_i s_{i+2} s_{i+1} \mathbf{y}_1 s_i \mathbf{y}_2 s_i s_{i+1} \mathbf{y}_4 s_i \mathbf{z}_2],$$

where $\mathbf{y}_4 = \mathbf{y}_3 \mathbf{z}_1$. The generator s_{i-1} must appear in $\mathbf{y}_1$, or the subword $[s_i s_{i+2} s_{i+1} \mathbf{y}_1 s_i]$ would contradict Lemma 4.3.2 (i). We can thus write $\mathbf{y}_1 = \mathbf{y}'_1 s_{i-1} \mathbf{y}''_1$, where s_{i-1} does not appear in $\mathbf{y}''_1$. A similar argument applied to the subword $[s_i s_{i+1} \mathbf{y}_4 s_i]$ shows that s_{i-1} must appear in $\mathbf{y}_4$, and we can write $\mathbf{y}_4 = \mathbf{y}'_4 s_{i-1} \mathbf{y}''_4$, where s_{i-1} does not appear in $\mathbf{y}'_4$. The trace is now equal to

$$[x s_i s_{i+2} s_{i+1} \mathbf{y}'_1 s_{i-1} \mathbf{y}''_1 s_i \mathbf{y}_2 s_i s_{i+1} \mathbf{y}'_4 s_{i-1} \mathbf{y}''_4 s_i \mathbf{z}_2].$$

Note that we have

$$[s_{i+1} \mathbf{y}'_1 s_{i-1} \mathbf{y}''_1 s_i] = [\mathbf{y}'_1 s_{i+1} s_{i-1} s_i \mathbf{y}''_1],$$

since $\mathbf{y}'_1$ has no occurrences of s_i or s_{i+2} . Recall that $\mathbf{y}'_4$ contains no occurrences of s_{i-1}, s_{i+1} or s_i . It follows that the trace $[s_i s_{i+1} \mathbf{y}'_4 s_{i-1}]$ is equal to a trace of the form $[\mathbf{y}'_5 s_i s_{i+1} s_{i-1} \mathbf{y}''_5]$, where the indicated occurrences of s_{i-1}, s_i and s_{i+1} are identified, $\mathbf{y}'_5$ contains no occurrences of s_j for $j > i - 2$, and $\mathbf{y}''_5$ contains no occurrence of s_j for $j < i + 2$. The inductive hypothesis can now be applied to the subword

$$[\mathbf{y}'_1 s_{i+1} s_{i-1} s_i \mathbf{y}''_1 \mathbf{y}_2 \mathbf{y}'_5 s_i s_{i+1} s_{i-1} \mathbf{y}''_5],$$

which completes the proof. ■

Lemma 4.3.5 *Let X be a Coxeter graph of type H_n and let E be the heap of $w \in W_c(X)$. Assume that $m(s_1, s_2) = 5$. Then E cannot have a trace of the form*

$$[x(s_i s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2} s_i) \mathbf{z}]$$

in which $\mathbf{y}$ has no occurrences of s_{i+2} or s_{i+3} .

Proof. Assume that $i = 1$. Then

$$[\mathbf{x}(s_i s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2} s_i) \mathbf{z}] = [\mathbf{x}(s_3 s_1 s_2) \mathbf{y}(s_2 s_1 s_3) \mathbf{z}]$$

where $s_3, s_4 \notin \mathbf{y}$. We may assume that $\mathbf{y}$ is as short as possible, which means that only s_1 and s_2 can appear in $\mathbf{y}$. Now, $\mathbf{y}$ is non-empty, else $[\mathbf{x}(s_3 s_1 s_2) \mathbf{y}(s_2 s_1 s_3) \mathbf{z}]$ would not be reduced. Since $s_1 s_2 \mathbf{y} s_2 s_1$ has length greater than 4 and $m(s_1, s_2) = 5$, we must have $\mathbf{y} = s_1$. But this contradicts Theorem 3.3.8 (i).

Assume now that $i > 1$. We may assume that $\mathbf{y}$ is as short as possible, which means that only s_j with $j \leq i + 1$ can appear in $\mathbf{y}$. Since $m(s_i, s_{i+1}) = 3$, we can now use the argument dealing with the case $i > 2$ from Lemma 4.3.4, and the result follows. $\blacksquare$

Lemma 4.3.6 *Let X be a Coxeter graph of type $\tilde{C}_n$, (n odd) and let E be the heap of $w \in W_c(X)$. Then E has no trace of the following forms:*

- (i) $[\mathbf{x}(s_{i+2} s_i s_{i+1}) \mathbf{y}(s_{i+1} s_i) \mathbf{z}]$ in which $\mathbf{y}$ has no occurrences of s_i or s_{i-1} ;
- (ii) $[\mathbf{x}(s_i s_{i+1}) \mathbf{y}(s_{i+1} s_i s_{i+2}) \mathbf{z}]$ in which $\mathbf{y}$ has no occurrences of s_i or s_{i-1} ;
- (iii) $[\mathbf{x}(s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2} s_i) \mathbf{z}]$ in which $\mathbf{y}$ has no occurrences of s_{i+2} or s_{i+3} ;
- (iv) $[\mathbf{x}(s_i s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2}) \mathbf{z}]$ in which $\mathbf{y}$ has no occurrences of s_{i+2} or s_{i+3} .

Proof. By left and right symmetry and symmetry of the Coxeter graph, it is enough to prove (iv). The proof is by induction on i . Assume $i = 1$. We have

$$[\mathbf{x}(s_i s_{i+2} s_{i+1}) \mathbf{y}(s_{i+1} s_{i+2}) \mathbf{z}] = [\mathbf{x}(s_3 s_1 s_2) \mathbf{y}(s_2 s_3) \mathbf{z}]$$

where there are no occurrences of s_3 or s_4 in $\mathbf{y}$. We may assume $\mathbf{y}$ is as short as possible, which means that only s_1 and s_2 can appear in $\mathbf{y}$. Now, $\mathbf{y}$ is non-empty, else $[\mathbf{x}(s_3 s_1 s_2) \mathbf{y}(s_2 s_3) \mathbf{z}]$ would not be reduced. For the same reason, $\mathbf{y} \neq s_2$. If $\mathbf{y} = s_1$, then Theorem 3.3.8 (i) is violated since $m(s_1, s_2) = 4$. Thus $s_1 s_2 \mathbf{y} s_2$ has length greater than 4, so $s_1 s_2 \mathbf{y} s_2$ is not reduced, a contradiction.

Assume now that $i > 1$. We can write the trace in the form

$$[\mathbf{x} s_i s_{i+2} s_{i+1} \mathbf{y}_1 s_i \mathbf{y}_2 s_i s_{i+1} \mathbf{y}_4 s_i \mathbf{z}_2],$$

where the notation is as in the proof of Lemma 4.3.4. The proof is completed by copying the steps of the argument used to deal with the term $\mathbf{y}_1$ in the case $i > 2$ in Lemma 4.3.4. $\blacksquare$

4.4 The non-acyclic case

Recall from Definition 3.2.3 that R is the relation defined on the vertices of a heap E given by $v_1 R v_2$ if and only if $\partial_E(e_0) = v_1 + v_2$ for some edge e_0 of E .

Lemma 4.4.1 *Let X be a Coxeter graph of type $\tilde{F}_5$ and let E be the heap of $w \in W_c(X)$. Let α be a boundary vertex in E . Then either α is an effective boundary vertex or there exists an edge e in E such that $\partial(e) = \alpha + \beta$, where β is an effective boundary vertex and $\varepsilon(\alpha) \neq \varepsilon(\beta)$.*

Proof. Let the vertices of X be labelled $s_0, s_1, \dots, s_5$ and assume that $m(s_2, s_3) = 4$. By Theorem 3.4.1, there is a sequence

$$\alpha_0 R \alpha_1 R \dots R \alpha_k$$

in which α_0 is effective. By hypothesis, we have $k > 0$. Since X is symmetric, we can deal with the case where $\varepsilon(\alpha_0) = s_2$. Since α_0 is effective, we have $\varepsilon(\alpha_0) = s_2$ or s_3 . If $\varepsilon(\alpha_0) = s_0$, we must have $k = 1$ since there is no generator to the left of s_0 , and we are done.

The only other possibility is that $\varepsilon(\alpha_1) = s_2$. Then E has a trace of the form

$$[xs_1s_2\mathbf{y}s_3s_2s_1s_3\mathbf{z}],$$

where $\mathbf{y}$ has no occurrences of s_0, s_1 or s_2 .

Suppose for a contradiction that $\mathbf{y}$ contains an occurrence of s_3 . Then the trace is equal to

$$[xs_1s_2\mathbf{y}_1s_3\mathbf{y}_2s_3s_2s_1\mathbf{z}],$$

where there are no occurrences of s_3 or s_2 in $\mathbf{y}_2$ or $\mathbf{y}_3$. Applying Lemma 4.3.2 (i) to $[s_3\mathbf{y}_2s_3]$ gives the required contradiction.

Since $\mathbf{y}$ has no occurrences of s_3 , we have $[s_2\mathbf{y}] = [\mathbf{y}s_2]$, and

$$[xs_1s_2\mathbf{y}s_3s_2s_1s_3\mathbf{z}] = [xs_1\mathbf{y}s_2s_3s_2s_1s_3\mathbf{z}] = [xs_1\mathbf{y}s_2s_3s_2s_3s_1\mathbf{z}],$$

which contradicts Theorem 3.3.8 (i). ■

Definition 4.4.2 Let E be a heap and let

$$c = \{\alpha_1, \alpha_2, \dots, \alpha_r\} : \alpha_1 < \alpha_2 < \dots < \alpha_r$$

be a chain in E . We say c is balanced if $\varepsilon(\alpha_1) = \varepsilon(\alpha_r)$. If c is a balanced convex chain, we define the heap $E \setminus c$ to be the subheap of E obtained by omitting the vertices $\alpha_2, \alpha_3, \dots, \alpha_r$. We call the heap $E \setminus c$ the contraction of E along c , and the number r the length of the chain.

Remark 4.4.3 If, in the above definition, we had omitted vertices $\alpha_1, \alpha_2, \dots, \alpha_{r-1}$ instead, we would have obtained the the same heap: the fact that c is balanced makes the two corresponding labelled heaps isomorphic.

Lemma 4.4.4 Let E be a heap and let c be a balanced convex chain of E .

(i) If the length of c is 2, then $\dim \ker \partial_E = \dim \ker \partial_{E \setminus c} + 1$.

(ii) If c is a balanced convex chain $x < y < z$ such that $\varepsilon(x) \neq \varepsilon(y)$, then $\dim \ker \partial_E = \dim \ker \partial_{E \setminus c}$.

Proof. Part (i) is [5, Lemma 2.3.4] and part (ii) is [5, Lemma 2.3.5]. ■

Lemma 4.4.5 Let $W = W(X)$ be a star reducible Coxeter group with simply laced or complete Coxeter graph X and let $\mathbf{s}$ be a reduced expression for $w \in W_c$. If $x = sw \notin W_c$, then the heap $E = \overline{\phi}(s[\mathbf{s}])$ is acyclic.

Proof. In the simply laced case, this is a restatement of [5, Theorem 2.4.2 (ii)], which shows that if $\overline{\phi}(s[\mathbf{s}])$ is not acyclic, then $[\mathbf{s}] = [\mathbf{s}\mathbf{u}]$ for some $\mathbf{u}$, in which case $x = sw \in W_c$. If X is complete, E is totally ordered and E will have Property P1 as long as $s[\mathbf{s}]$ has no trace of the form $[xs_i s_i \mathbf{y}]$. This is true since $s[\mathbf{s}]$ is reduced. ■

Lemma 4.4.6 *Let $W(X)$ be a star reducible Coxeter group, let s be a reduced expression for $w \in W_c$ and let $x = sw \notin W_c$ for some $s \in S$. Let E be the heap of w and assume $\overline{\phi}(s[s])$ is not acyclic. If a boundary vertex α occurs in a sequence $\alpha_0 R \alpha_1 R \cdots R \alpha_k = \alpha$ where $\alpha_0, \alpha_1, \dots, \alpha_k = \alpha$ are boundary vertices and α_0 is an effective boundary vertex, then $\varepsilon(\alpha_i) \neq \varepsilon(\alpha_{i+1})$, except possibly if $i = k - 1$.*

Proof. The simply laced and complete Coxeter graphs are dealt with in Lemma 4.4.5. Thus X is a Coxeter graph of type $B_n, F_n, \tilde{C}_n, (n \text{ odd}), H_n$, or $\tilde{F}_5$. This means that if $e_1 = (x_1, y_1)$ and $e_2 = (x_2, y_2)$ are edges in E with $\partial(e_1) = \alpha_l + \alpha_{l+1}$ and $\partial(e_2) = \alpha_{l+1} + \alpha_{l+2}$, then the subscripts of the Coxeter generators $\varepsilon(x_1), \varepsilon(x_2)$ differ by 0 or 2.

Suppose we are in type $\tilde{C}_n, (n \text{ odd})$, and let l be minimal such that $\varepsilon(\alpha_l) = \varepsilon(\alpha_{l+1})$ as in the statement. Let e be an edge of E such that $\partial(e) = \alpha_l + \alpha_{l+1}$. By symmetry, we may assume that $e = (x, y)$ with $\varepsilon(x) = \varepsilon(y) = s_{j-1}$. This cannot happen without contradicting Lemma 4.3.6 (i) or (iii) with $i = j - 1$. This completes the proof in type $\tilde{C}_n, (n \text{ odd})$, which contains type B_n as a subcase.

The case where X is a Coxeter graph of type $\tilde{F}_5$ is dealt with in Lemma 4.4.1, which shows that $\varepsilon(\alpha_i) = \varepsilon(\alpha_{i+1})$ is impossible. Thus the present argument reduces to types F_n and H_n .

Case (i). Let X be a Coxeter graph of type F_n . Suppose for a contradiction that we have $\alpha_l R \alpha_{l+1} R \alpha_{l+2}$ with $\varepsilon(\alpha_l) = \varepsilon(\alpha_{l+1}) = \varepsilon(\alpha_{l+2}) = s_j$. Then there are edges $e_1 = (x_1, y_1)$ and $e_2 = (x_2, y_2)$ in E with $\partial(e_1) = \alpha_l + \alpha_{l+1}$ and $\partial(e_2) = \alpha_{l+1} + \alpha_{l+2}$. We must have $\varepsilon(x_1) = s_{j-1}$ and $\varepsilon(x_2) = s_{j+1}$ or vice versa. We deal with the first case; the other case is symmetric. The trace of E is now of the form

$$[x s_{j+1} s_j \mathbf{y}_1 s_{j-1} s_j s_{j+1} \mathbf{y}_2 s_j s_{j-1} \mathbf{z}]$$

or

$$[\mathbf{z} s_{j-1} s_j \mathbf{y}_2 s_{j+1} s_j s_{j-1} \mathbf{y}_1 s_j s_{j+1} \mathbf{x}],$$

in which $\mathbf{y}_1$ has no occurrences of s_j or s_{j+1} and $\mathbf{y}_2$ has no occurrences of s_j or s_{j-1} .

Let t be the unique index with $m(s_t, s_{t+1}) > 3$. If $t \geq j$, then the subword $[s_j \mathbf{y}_1 s_{j-1} s_j]$ or $[s_j s_{j-1} \mathbf{y}_1 s_j]$ contains no occurrences of s_{j+1} , which contradicts Lemma 4.3.2 (ii). If $t < j$, then the subword $[s_j s_{j+1} \mathbf{y}_2 s_j]$ or $[s_j \mathbf{y}_2 s_{j+1} s_j]$ contradicts Lemma 4.3.2 (i).

Now suppose $\varepsilon(\alpha_1) = \varepsilon(\alpha_0) = s_j$. If $j = 2$, this contradicts Lemma 4.3.3 (ii) or (iv); if $j = 3$, this contradicts Lemma 4.3.3 (iii) or (v). It is now enough to show we cannot have $s_j = \varepsilon(\alpha_l) = \varepsilon(\alpha_{l+1}) \neq \varepsilon(\alpha_{l+2}) = \varepsilon(\alpha_{l-1}) = s_{j'}$.

Assume first that $j' = j + 2$. In this case the trace is of the form

$$[x s_{j-1} s_{j+1} s_j s_{j+2} s_{j+1} \mathbf{y} s_{j+1} s_j s_{j+2} s_{j-1} s_{j+1} \mathbf{z}]$$

where $\mathbf{y}$ has no occurrences of s_{j-1} or s_j . Now the subword $s_j s_{j+2} s_{j+1} \mathbf{y} s_{j+1} s_j$ contradicts Lemma 4.3.2 (i) unless $j = 2$, in which case we contradict Lemma 4.3.3 (ii) or (iv).

We now deal with the case $j' = j - 2$. There exist distinct edges $e_1 = (x_1, y_1)$, $e_2 = (x_2, y_2)$, and $e_3 = (x_3, y_3)$ in E with $\partial(e_1) = \alpha_{l-1} + \alpha_l$, $\partial(e_2) = \alpha_{l+1} + \alpha_l$, and $\partial(e_3) = \alpha_{l+1} + \alpha_{l+2}$. Assume without loss of generality that $x_3 < y_1$. The formula for $\partial(e_3)$ shows that the trace of E has subwords of the form $s_{j-2} s_j s_{j-1}$ and $s_{j-1} s_j s_{j-2}$, where the first subword occurs to the left of the second subword. If the occurrences of s_{j-1} are the same, the trace has a subword of the form

$$[s_{j-2} s_j s_{j-1} s_j s_{j-2}] = [s_j s_{j-2} s_{j-1} s_{j-2} s_j],$$

which is impossible as either $m(s_{j-2}, s_{j-1}) = 3$ or $m(s_j, s_{j-1}) = 3$. If the occurrences are distinct, then the trace is of the form

$$[\mathbf{x}s_{j-1}s_{j+1}s_{j-2}s_{j-1}\mathbf{y}s_{j-1}s_{j-2}s_{j+1}s_{j-1}\mathbf{z}]$$

where $\mathbf{y}$ has no occurrences of s_j or s_{j+1} , which contradicts Lemma 4.3.4 with $i = j - 2$. This completes the proof in type F_n .

Case (ii). Let X be a Coxeter graph of type H_n and let $s_j = \varepsilon(\alpha_0)$; note that $j = 1$ or 2 . If $\varepsilon(\alpha_1) = \varepsilon(\alpha_0) = s_j$, then we cannot have $j = 1$ because s_1 is the leftmost vertex. If $j = 2$, then we must have $\partial(e) = \alpha_0 + \alpha_1$ for some edge $e = (x, y)$ with $\varepsilon(x) = s_3$. If $k > 1$, we must have $\varepsilon(\alpha_2) = s_1$ (as opposed to s_3). This implies that $\varepsilon(\alpha_0) = \varepsilon(\alpha_1) = \varepsilon(\alpha_2)$, which was proved impossible above.

We may therefore assume that $\varepsilon(\alpha_0) \neq \varepsilon(\alpha_1)$. It is now enough to show that we cannot have

$$s_j = \varepsilon(\alpha_l) = \varepsilon(\alpha_{l+1}) \neq \varepsilon(\alpha_{l+2}) = \varepsilon(\alpha_{l-1}) = s_{j'}.$$

We now argue as in Case (i) above, except at the final step, where we invoke Lemma 4.3.5 instead of Lemma 4.3.4. (In this case, the situation where $j = 2$ in Lemma 4.3.3 cannot arise.) ■

Lemma 4.4.7 *Let $W(X)$ be a star reducible Coxeter group, let $\mathbf{s}$ be a reduced expression for $w \in W_c$ and let $x = sw \notin W_c$ for some $s \in S$. Let E be the heap of w and assume $\bar{\phi}(s[\mathbf{s}])$ is not acyclic. Let $\mathbf{c} = s^{(0)} < t^{(1)} < s^{(1)} < t^{(2)} < \dots < s^{(k)}$ (or $t^{(k)}$, depending on the parity of $m(s, t)$) be the convex chain in $s[\mathbf{s}]$ described in Theorem 3.3.8 (i), where $s[\mathbf{s}] = [\mathbf{x}'\mathbf{c}\mathbf{x}'']$ and $\mathbf{c}' = \mathbf{c} \setminus \{\alpha_r\}$. Then the heap $\bar{\phi}(\mathbf{x}'\mathbf{c}'\mathbf{x}'')$ is acyclic.*

Proof. If X is simply laced or complete, then the heap $\bar{\phi}(s[\mathbf{s}])$ is acyclic by Lemma 4.4.5, so we may assume X is a straight line graph. Let $E_0 = \bar{\phi}(s[\mathbf{s}])$ and let $s^{(0)}$ denote the new vertex in E_0 . Since E is acyclic by Lemma 3.3.18 (i) (iv), Lemma 4.1.1 shows that $\dim \ker \partial_{E_0} = 1$. Now $e_0 = (s^{(0)}, s^{(1)})$ is an edge such that $\partial(e_0) = t^{(1)}$. Since E_0 is not acyclic, there exist edges $e_0, e_1, \dots, e_r$ in E_0 and scalars $\lambda_0, \lambda_1, \dots, \lambda_r$, not all equal to 0, such that

$$\partial_{E_0} \left(\sum_{i=0}^r \lambda_i e_i \right) = 0,$$

where $e_0 = (s^{(0)}, s^{(1)})$, so $\partial_E(e_0) = t^{(1)}$. Since E is acyclic and E is a convex subheap of E_0 by Lemma 3.3.9, Lemma 3.3.12 shows that $\lambda_0 \neq 0$, and without loss of generality we may assume $\lambda_0 = 1$. Now we have that

$$\partial_{E_0} \left(- \sum_{i=1}^r \lambda_i e_i \right) = t^{(1)}$$

and similarly,

$$\partial_E \left(- \sum_{i=1}^r \lambda_i e_i \right) = t^{(1)}$$

by Lemma 3.3.12. Therefore $t^{(1)}$ is a boundary vertex in E .

By Theorem 3.4.1, there is a chain of linear equivalences $\alpha_0 R \alpha_1 R \dots R \alpha_k$, where $\alpha_k = t^{(1)}$, $\alpha_0, \alpha_1, \dots, \alpha_k$ are boundary vertices in E , and α_0 is effective. By Lemma 4.4.6, the set $L = \{\varepsilon(\alpha_0), \varepsilon(\alpha_1), \dots, \varepsilon(\alpha_k)\}$ has cardinality $k+1$ or k , and in the latter case we have $\varepsilon(\alpha_{k-1}) = \varepsilon(\alpha_k)$. The proof of Lemma 4.4.6 shows that the latter situation can only occur in type F_n or H_n .

Case (i). Suppose first that $|L| = k + 1$. In this case, $\alpha_0, \alpha_1, \dots, \alpha_k$ correspond to the labels $s_2, s_4, \dots, s_{k+2}$ in Figure 8 below; this implies that $m(s, t) = 3$.




Figure 8

By Lemma 4.4.4, if we contract the chain $c = s_1 < s_2 < s_1$, then $\dim \ker \partial_{E_0} = \dim \ker \partial_{E_0 \setminus c} = 1$. If we repeat this process by contracting the chain $c' = s_3 < s_4 < s_3$ in $E_0 \setminus c$, then $\dim \ker \partial_{E_0 \setminus c} = \dim \ker \partial_{(E_0 \setminus c) \setminus c'} = 1$. By iterating this process, we wind up with the heap E' in Figure 9 below.



Figure 9

Contracting along $c'' = s_{2k+3} < s_{2k+3}$ then gives

$$\dim \ker \partial_{E'_0 \setminus c''} = \dim \ker \partial_{E'_0} - 1.$$

Since $\dim \ker \partial_{E_0} = 1$, $E'_0 \setminus c''$ is acyclic. Since the lower occurrence, $\gamma = s^{(1)}$, of s_{2k+3} is only involved in the last contraction, $\dim \ker \partial_{E_0 \setminus s^{(1)}} < \dim \ker \partial_{E_0} = 1$. Thus $E_0 \setminus s^{(1)}$ is acyclic, as required.

Case (ii). Now assume $|L| = k$ and X is of type H_n or F_n . Assume first that the chain $\alpha_0 R \alpha_1 R \dots R \alpha_k = \alpha$ and its corresponding edges is as shown up to symmetry in Figure 10 below, where the four occurrences of s_{k-1} are distinct and $\varepsilon(\alpha_l) = s_{2l+2}$ for $l < k$; this implies that $m(s, t) = 3$. We proceed as in case (i) starting with the chain $s_1 < s_2 < s_1$, removing the α_i in order starting with α_0 and ending with the chain $s^{(0)} < s^{(1)}$. Since $s^{(1)}$ is not involved with any earlier contractions, we argue as in case (i) that $E_0 \setminus s^{(1)}$ is acyclic.




Figure 10

Case (iii). The other possibility is that $|L| = k$ and X is of type H_n or F_n , but the occurrences of s_{k-1} in Figure 10 are the same. This situation is shown in Figure 11 below.




Figure 11

In this case, the occurrences of s_k shown are $t^{(1)}$ and $t^{(2)}$, which implies that $m(s_{k-1}, s_k) > 3$. Since α_0 is effective, $m(s_1, s_2) > 3$. The only way this can happen is if $k = 2$ and X is of type H_n . This situation is shown in Figure 12, where the bottom left s_1 is $s^{(2)}$ and $m(s, t) = 5$.




Figure 12

In this case, we perform a contraction along $\mathbf{c} = s^{(0)} < t^{(1)} < s^{(1)}$, followed by a contraction along $\mathbf{c}' = s_3 < s_2 < s_3$. Since $s^{(2)}$ is not involved in either contraction, we can argue as before to show that $\dim \ker \partial_{E_0 \setminus s^{(2)}} = 0$, which completes the proof. ■

Theorem 4.4.8 (Property W) *Let $W(X)$ be a star reducible Coxeter group with Coxeter graph X . Let $w \in W_c$ and let $x = sw \notin W_c$ for some $s \in S$. Then $\tilde{t}_x \in v^{-1}\mathcal{L}$.*

Proof.

Let E be the heap of w and let $[s]$ be the trace of E . If $\bar{\phi}(s[s])$ is acyclic, the conclusion follows by Lemma 4.2.1. In particular, if X is simply laced or complete we are done by Lemma 4.4.5, so we may assume from now on that $\bar{\phi}(s[s])$ is not acyclic.

Assume that $x = sw = u'w_{st}u''$ reduced, where w_{st} is the longest element in the parabolic subgroup $\langle s, t \rangle$. We can use the relation

$$\tilde{t}_{w_{st}} = - \sum_{\substack{u \in \langle s, t \rangle \\ u < w_{st}}} v^{\ell(u) - m(s, t)} \tilde{t}_u$$

to write

$$\begin{aligned} \tilde{t}_x &= \tilde{t}_{u'} \tilde{t}_{w_{st}} \tilde{t}_{u''} \\ &= -\tilde{t}_{u'} \left(v^{-1} \tilde{t}_{tw_{st}} + v^{-1} \sum_{u \leq sw_{st}} v^{\ell(u) - \ell(sw_{st})} \tilde{t}_u \right) \tilde{t}_{u''}. \end{aligned}$$

Arguing as in Case (v) of the proof of Lemma 4.2.1, we find that all of the terms in the expression for $\tilde{t}_x$ arising from $u \leq sw_{st}$ lie in $v^{-1}\mathcal{L}$. Now write $s[s] = \mathbf{x}'\mathbf{c}\mathbf{x}''$ and $\mathbf{x}'\mathbf{c}'\mathbf{x}''$ as in Lemma 4.4.7. By Lemma 4.4.7, $\bar{\phi}([\mathbf{x}'\mathbf{c}'\mathbf{x}''])$ is acyclic. By Lemma 4.1.6, $\tilde{t}_{\mathbf{x}'\mathbf{c}'\mathbf{x}''} \in \mathcal{L}$. Now,

$$\tilde{t}_{\mathbf{x}'\mathbf{c}'\mathbf{x}''} = \tilde{t}_{u'} \tilde{t}_{tw_{st}} \tilde{t}_{u''},$$

so $v^{-1} \tilde{t}_u \tilde{t}_{tw_{st}} \tilde{t}_{u''} \in v^{-1}\mathcal{L}$, which completes the proof. ■

Bibliography

- [1] M. Baker and S. Norine. Riemann–Roch and Abel–Jacobi theory on a finite graph. *Adv. Math.*, 215:766–788, 2007.
- [2] A. Björner and F. Brenti. *Combinatorics of Coxeter Groups*. Springer-Verlag, 2002.
- [3] M. Geck and G. Pfeiffer. *Characters of Finite Coxeter Groups and Iwahori–Hecke Algebras*. Oxford University Press, 2000.
- [4] J.J. Graham. *Modular representations of Hecke algebras and related algebras*. PhD thesis, University of Sydney, 1995.
- [5] R.M. Green. Acyclic heaps of pieces, I. *J. Algebraic Combin.*, 19:173–196, 2004.
- [6] R.M. Green. Star reducible Coxeter groups. *Glasgow Math. J.*, 48:583–609, 2006.
- [7] R.M. Green. Generalized Jones traces and Kazhdan–Lusztig bases. *J. Pure Appl. Alg.*, 211:744–772, 2007.
- [8] R.M. Green and J. Losonczy. Canonical bases for Hecke algebra quotients. *Math. Res. Lett.*, 6:213–222, 1999.
- [9] J.E. Humphreys. *Reflection Groups and Coxeter Groups*. Cambridge University Press, 1990.
- [10] D. Kazhdan and G. Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.*, 53:165–184, 1979.
- [11] G. Lusztig. Cells in affine Weyl groups. *J. Fac. Sci. Tokyo U.*, I:297–328, 1985.
- [12] J.R. Stembridge. On the fully commutative elements of Coxeter groups. *J. Algebraic Combin.*, 5:353–385, 1996.
- [13] G.X. Viennot. Heaps of pieces, I: basic definitions and combinatorial lemmas. In ed. G. Labelle and P. Leroux, editors, *Combinatoire Énumérative*, pages 321–350. Springer-Verlag, 1986.